OFFICIAL

Algal bloom wildlife post-mortem report



Species - Pied cormorant

Date collected - 25 June 2025

Location - Bay of Shoals, Kangaroo Island

History relating to the animal

Eight pied cormorants (*Phalacrocorax varius*) were submitted from a single mass mortality event from Bay of Shoals, Kangaroo Island.

Examination

Three birds were seen alive, weak with paralysis (weakness of limbs) and ataxia (wobbly). Two of these birds were on the sandy beach and were unable to fly and ataxic (wobbly). The third bird was in the shallow water and was unable to swim and circling (moving in circles with no apparent intent). The affected birds were euthanised on humane grounds. Another 5 deceased birds were found in the same area and were submitted to the laboratory. These animals were already dead and so could not be examined prior to death.

Necropsy

The necropsies (looking at the whole body) revealed that the animals had mild effects from decomposition. The birds were in moderate to poor body condition with atrophy (wasting) of muscle over the pectoral (breast) bone. There was no food present in the proventriculus/ventriculus (the top part of a bird's stomach/ stomach). There were nematodes (round worms) present in the gastrointestinal tract (gut), which is a normal finding in wild cormorants. All birds were adults, of mixed sex, except for one which was a juvenile female.

Tissues were collected to test for avian influenza testing and Newcastle disease, and for histopathology (looking at tissues under the microscope for more detailed information). Testing for brevetoxins and other algal biotoxins, a possibility due to the algal bloom, was requested. Samples were collected for other testing that might be indicated after these tests were performed.

Histopathology

Samples from every major body system were examined under the microscope for the three freshest birds. All organs appeared normal (including the presence of the nematodes) except for the kidney and liver in bird 2.

In the kidney, there were ova (eggs) of a parasite (fluke worm) that was an incidental finding and has previously been found in cormorants.

In the liver there was a bacterial infection that was causing tissue necrosis (tissue death).

The diagnosis from the histopathology was that the hepatitis contributed to the death in bird 2.

OFFICIAL

Algal bloom wildlife post-mortem report



Culture

Following the histopathology findings, which suggested a possible infection, the liver of bird 2 and 3 were cultured to try to identify bacteria.

In bird 2, there was no bacterial growth after 48 hours, and anaerobes (those bacteria that do not need an oxygenated environment), including *Salmonella* species, *Listeria* species and *Yersina* species, were not isolated.

In bird 3, there was a light growth of *Clostridium perfringens* (a common, widespread bacteria) and mixed organisms (e.g. bacteria), but anaerobes including *Salmonella* species, *Listeria species* and *Yersina* species, were not isolated. The bacteria found were not clinically significant (not likely to be causing disease).

Brevetoxins

Brevetoxin 3 was found in the brain (0.01 mg/kg), liver (0.09 mg/kg) and lung (0.02 mg/kg) of bird 1, in the brain (0.02 mg/kg), liver (0.03 mg/kg) and lung (0.01 mg/kg) of bird 2 and the lung (0.01 mg/kg) of bird 3.

Other algal biotoxins

Results were below reporting limits or the sample was insufficient for testing.

Avian influenza

Results were negative.

Newcastle disease

Results were negative.

Summary

Eight pied cormorants were found dead or in poor condition at Bay of Shoals, Kangaroo Island.

Of these, three live birds were observed with neurological symptoms (weakness of limbs, wobbly). The remaining five birds were already dead.

Laboratory examination (necropsy and histopathology) on all birds could not determine the cause of chronic weight loss, neurological signs and death. Hepatitis (liver inflammation) may have contributed to one of the birds being unwell.

Three birds were sent for additional testing.

Testing for avian influenza and Newcastle disease was negative.

OFFICIAL

Algal bloom wildlife post-mortem report



Brevetoxins were detected in the tissues of all birds tested. The levels detected are consistent with those found in marine birds overseas that were reported to have been exposed to harmful algal blooms.

It cannot be determined whether the brevetoxin presence in the cormorants was a direct contributor to their death or a non-lethal co-morbidity given the poor condition of the birds at death. Pied cormorants primarily feed on fish. It is considered that the most significant exposure of the cormorants to brevetoxins is likely to have been via their consumption of fish organs containing brevetoxins.





Referred on 07/07/25 by

Owner:

COMORANT MORTALITY KANGAROO ISLAND SHAOLS BEACH Animal/s:

Wild Birds

DOB: N/A

Collected: 07/07/25 00:25 Subm.No:

Lab No.:

Samples tested as received

NECROPSY REPORT

CLINICAL HISTORY

Please refer to the clinical history on the request form and the clinical notes sent with the request form. A brief summary of the clinical history;

Dead cormorants and cormorants with neurological signs were found on Shoal Beach, Kangaroo Island

SAMPLES SUBMITTED

8 dead cormorants (Phalacrocorax sp.)
The birds are allocated numbers 1-8 at the laboratory

NECROPSY FINDINGS

These descriptions refer to all birds;

There is atrophy of the pectoral muscles. There is no ingesta in the proventriculus / ventriculus. There are low numbers of ascarids $30-50\,\mathrm{mm}$ long in the proventriculus / ventriculus and intestines. All birds are adults except for bird 8.

Additional findings

1

The tag on the bag is " this is the freshest bird"
This is a male bird. The bird is in moderate to poor body condition,
there is atrophy of both pectoral muscles and the bird weighs 1.1 kg.
There are minimal post mortem autolytic changes.

- 2 This is a male bird. The bird is in moderate to poor body condition, there is atrophy of both pectoral muscles and the bird weighs 1.0 kg. There are minimal post mortem autolytic changes.
- 3
 This is a female bird. The bird is in moderate to poor body condition, there is atrophy of both pectoral muscles and the bird weighs 1.2 kg

There are minimal post mortem autolytic changes.

This is a female bird. The bird is in moderate to poor body condition, there is atrophy of both pectoral muscles and the bird weighs 1.5 kg There are mild post mortem autolytic changes.





Neteried on 07707723 by

Owner:

COMORANT MORTALITY KANGAROO ISLAND SHAOLS BEACH Animal/s:

Wild Birds

DOB: N/A

Collected: 07/07/25 00:25 **Subm.No:**

Lab No.:

Samples tested as received

5

This is a male bird. The bird is in moderate to poor body condition, there is atrophy of both pectoral muscles and the bird weighs 1.0 kg There are moderate post mortem autolytic changes.

This is a female bird. The bird is in moderate to poor body condition, there is atrophy of both pectoral muscles and the bird weighs 1.0 kg There are marked post mortem autolytic changes and maggots in the oral cavity.

This is a female bird. The bird is in moderate to poor body condition, there is atrophy of both pectoral muscles and the bird weighs 1.0 kg There are moderate post mortem autolytic changes.

This is a female juvenile bird. The bird is in moderate to poor body condition, there is atrophy of both pectoral muscles and the bird weighs $0.5\ kg$ There are moderate post mortem autolytic changes.

GROSS SUMMARY Birds 1-8 Chronic weight loss Alimentary ascardiasis

SAMPLES COLLECTED & TESTING

Birds 1-3

Cloacal and tracheal swabs are pooled for 1 x NDV PCR and 1 x AI PCR. Formalin fixed tissues are processed for histopathology. Fresh liver (10g), spleen, heart, lung, kidney, brain (< 500mg for these tissues) are stored frozen.

No samples are collected from birds 4-8

COMMENTS

The cause of chronic weight loss and death is not determined by gross findings. Alimentary ascaridasis (likely Anisakis or Contracaecum sp.) is a normal finding in cormorants.





Tested on 09/07/25

Reported on 13/08/25 17:30 **Referred on** 07/07/25 **by:**

Referred on 07/07/25 by:

Owner:

COMORANT MORTALITY KANGAROO ISLAND SHAOLS BEACH Animal/s:

Wild Birds

DOB: N/A

Collected: 07/07/25 00:25 Subm.No:

Lab No.:

Samples tested as received

Specialist Veterinary Anatomic Pathologist

Specialist Veterinary Anatomic Pathologist

Validated by



PATH RESULTS: COMORANT MORTALITY, (Wi)

From

Date Wed 13/08/2025 5:00 PM

To

Owner: COMORANT MORTALITY KANGAROO ISLAND SHAOLS BEACH

Collected: 07/07/25 00:25 Subm.No:

Soffected. 07707723 00.23 Subin. N

HISTOPATHOLOGY FROM NECROPSY

Tested on

Animal/s:

DOB: N/A

Wild Birds

Referred on

09/07/25

Lab No.:

07/07/25 **by**:

Reported on 13/08/25 17:30

REF: 2025/V HI 1249 CLINICAL HISTORY

Dead cormorants and cormorants with neurological signs were found on

Shoal Beach, Kangaroo Island

Samples tested as received

MACROSCOPY

Slides 1A-C to 3A-C correspond to birds 1-3; tissues for each bird include liver, spleen, heart, lung, kidney, brain, crop, proventriculus / ventriculus, duodenum, pancreas, jejunum, ileum, caecum

MICROSCOPY

Bird 1

Crop and proventriculus: There are multiple cross sections of ascarids (30-40 micron across, smooth cuticle, platymyarian musculature, pseudocoelom, lateral cords and alimentary tract)

There is mild autolysis of the alimentary sections.

Bird 2

Kidney: Multifocally the renal pelvises are expanded by abundant oval

ova (10 micron diameter with smooth capsule and operculum) within their lumens.

Liver: Multifocally necrotizing and heterophilic hepatitis with intralesional short rod bacteria effaced the hepatic acinar architecture. (Moderate, multifocal, subacute necrotizing heterophilic hepatitis with intralesional short rod bacteria)

There is proventricular ascaridiasis.

Bird 3

There is proventricular ascaridiasis.

Those tissues not described for each bird appear normal.

DIAGNOSIS Bird 1-3 Alimentary ascardiasis

Bird 2





Referred on 07707723 B

Owner:

COMORANT MORTALITY KANGAROO ISLAND SHAOLS BEACH Animal/s:

Wild Birds

DOB: N/A

Collected: 07/07/25 00:25 **Subm.No:**

Lab No.:

Samples tested as received

Renal trematode ova Bacterial hepatitis

COMMENTS

Bacterial hepatitis contributed to morbidity for bird 2. Please contact the laboratory within the next 5 working days if microbiological culture of bird 2 is required.

Alimentary ascardiasis (Anisakis sp. or Contracaecum sp.) are a normal finding in marine birds.

The trematode Renicola has been recorded in the kidneys of cormorants (P. Ladds Pathology of Australian native wildlife 2009). This is an incidental finding.

Specialist Veterinary Anatomic Pathologist

Validated by

MICROBIOLOGY

SPECIMEN: Liver
ANIMAL ID:BIRD 3

MICROSCOPY

No bacteria seen.

CULTURE

1. Light growth of Mixed organisms

COMMENT: Culture is continuing.

INTERIM REPORT
12/08/25
Validated by Laboratory Scientist.



Owner:

COMORANT MORTALITY KANGAROO ISLAND SHAOLS BEACH Animal/s:
 Wild Birds

WIIG BIIGS

DOB: N/A

Collected: 07/07/25 00:25 Subm.No:

Lab No.:

Samples tested as received

MICROBIOLOGY SPECIMEN: Liver ANIMAL ID:BIRD 2

MICROSCOPY

No bacteria seen.

A small number of leucocytes. Scanty epithelial cells.

CULTURE

1. No growth after 48 hours incubation.

COMMENT: Anaerobes NOT isolated.

No Salmonella sp., Listeria sp. or Yersinia sp. isolated.

Final Report
21/07/25
Validated by Laboratory Scientist.

Number of samples

9



Owner:

COMORANT MORTALITY
KANGAROO ISLAND
SHAOLS BEACH

Animal/s:
 Wild Birds

DOB: N/A

Collected: 07/07/25 00:25 Subm.No:

Lab No.:

Samples tested as received

MOLECULAR DIAGNOSTICS

NEWCASTLE DISEASE VIRUS RNA PCR (REAL TIME REVERSE TRANSCRIPTASE)

Specimen type: Tracheal + Number of specimens: 1

cloacal swabs in VTM

SPECIMEN ID F Gene M Gene L Gene

CORMORAN Not detected Not detected Not detected

Validated by , Laboratory Scientist.





Neteried on Otyoty23

Owner:

COMORANT MORTALITY KANGAROO ISLAND SHAOLS BEACH Animal/s:
 Wild Birds

DOB: N/A

Collected: 07/07/25 00:25 **Subm.No:**

Lab No.:

Samples tested as received

MOLECULAR DIAGNOSTICS

INFLUENZA A RNA PCR (REAL TIME REVERSE TRANSCRIPTASE)

Specimen type: Tracheal + cloacal swabs in VTM

SPECIMEN ID Type A H5 H7

CORMORAN Not detected

Validated by Laboratory Scientist.

CASE MANAGEMENT DETAILS

Case Managed by:

Case Management Requested by: Case Management Requested on:



Case Details:

Mortality



PATH RESULTS: COMORANT MORTALITY, (Wi)

From

Date Thu 14/08/2025 11:00 AM

То

Owner: COMORANT MORTALITY KANGAROO ISLAND

SHAOLS BEACH

Collected: 07/07/25 00:25 Subm.No:

Samples tested as received

MICROSCOPY

No bacteria seen.

CULTURE

MICROBIOLOGY

1. Light growth of Clostridium perfringens

2. Light growth of Mixed organisms

COMMENT: No Salmonella sp., Listeria sp. or Yersinia sp. isolated.

Final Report_

14/08/25 Validated by

Laboratory Scientist.

Tested on

09/07/25

Reported on 14/08/25 11:30

Referred on 07/07/25 by:

Animal/s: Wild Birds

DOB: N/A

Lab No.:

SPECIMEN: Liver ANIMAL ID:BIRD 3



PATH RESULTS: COMORANT MORTALITY, (Wi)

From

Date Tue 26/08/2025 11:00 AM



Owner: COMORANT MORTALITY KANGAROO ISLAND

SHAOLS BEACH

Collected: 07/07/25 00:25 Subm.No:

Samples tested as received

Tested on 09/07/25 Reported on 26/08/25 11:30

Referred on 07/07/25 **by**:

Animal/s: Wild Birds

DOB: N/A

Lab No.:

All Tests Complete

SUMMARY DIAGNOSIS Bird 2 Chronic hepatitis Renal trematodiasis

Birds 1-8 Alimentary ascardiasis

SUMMARY COMMENTS

The cause of chronic weight loss, neurological signs and death is not determined by laboratory tests. Avian influenza and Newcastle disease are excluded as causes of morbidity, neurological signs and death based on the negative PCR results.

Hepatitis contributed to morbidity for bird 2. Bacterial cultures are unremarkable. The renal trematode Renicola has been recorded in the kidneys of cormorants (P. Ladds 2009; Pathology of Australian native wildlife). This is an incidental finding.

The Cl. perfringens from bird 3 liver culture is a post mortem overgrowth.

Alimentary ascardiasis (Anisakis sp. or Contracaecum sp.) is a normal finding in marine birds.

Specialist Veterinary Anatomic Pathologist

Validated by



ANALYTICAL SERVICES TASMANIA

3 18 St Johns Avenue New Town 7008 TAS

C 03 6165 3300

enquiries@ast.tas.gov.au

www.analyticalservices.tas.gov.au

Submission Number: Report Number: Issue Date: Status:

4/09/2025 Final

CERTIFICATE OF ANALYSIS

Customer:
Address:
Contact:

Submission Description: Sample Received Date: Contract Number: Client Order Number: Program/Quote Reference: Cormorants Birds 1-3, Shoals Beach
22/08/2025

- Brevitoxin, lipophilic toxin and paralytic shellfish toxins

Sample(s) analysed as received. Sampling date and time data supplied by the client. The document shall not be reproduced except in full.

Additional information relating to this submission can be found in the sample receipt notification.

This report supersedes any previous reports with this submission number.

Many tests specify a holding time which gives the recommended timeframe by which a sample should be preserved/extracted and/or analysed after the sample is taken.

Holding time information can be found on the AST website https://analyticalservices.tas.gov.au/our-services/containers-samples-and-submissions.

Whilst every effort is made to analyse samples within these timeframes, situations can occur where this is not possible.

Where a test has been conducted outside the recommended sample holding time this should be taken into account when interpreting results.

The results in this report were authorised by:

Name Position

Section Head - Organic
Chemistry

Test Information:

3411 Lipophilic Toxins in Shellfish by LC-MS/MS 03-09-2025 3411A Brevetoxins in Biota by LC-MS/MS 02-09-2025	Method ID	Test Description	Date Commenced:
·	3411	Lipophilic Toxins in Shellfish by LC-MS/MS	03-09-2025
	3411A	Brevetoxins in Biota by LC-MS/MS	02-09-2025
3416 PST in Biota by LC-MS/MS (Boundy Method) 03-09-2025	3416	PST in Biota by LC-MS/MS (Boundy Method)	03-09-2025



0 1 0 1	
Sample Comments	
Sample Number:	
	3411A Brevetoxins in Biota by LC-MS/MS
	Brevetoxin 3
	The accuracy of this result may be affected by the small sample size provided.
Sample Number:	
IS - Insufficient Sam	pple. Sample not submitted.
Sample Number:	
	3411A Brevetoxins in Biota by LC-MS/MS
	Brevetoxin 3
	The accuracy of this result may be affected by the small sample size provided.
Sample Number:	
IS - Insufficient Sam	pple. Sample not submitted.
Sample Number:	
IS - Insufficient Sam	pple. Sample not submitted.
Sample Number:	
IS - Insufficient Sam	pple. Sample not submitted.

^{*}IS*- Insufficient Sample

^{*} NATA accreditation does not cover this result

Chemistry Test Results (Biota - Food)		Sample Description	Cormorant 1 Brain	Cormorant 1 Kidney	Cormorant 1	Cormorant 1 Lung	Cormorant 2 Brain 07/07/25 0:00	Cormorant 2 Kidney 07/07/25 0:00	Cormorant 2 Liver 07/07/25 0:00	Cormorant 2 Lung 07/07/25 0:00
		Sampled Date/ Time	07/07/25 0:00	07/07/25 0:00	07/07/25 0:00	07/07/25 0:00				
Method ID	Analyte	Units	338829	338830	338831	338832	338833	338834	338835	338836
	AZA1	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	AZA2	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	AZA3	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	< 0.01	<0.01
	Domoic Acid	mg/kg WMB	*IS*	*IS*	<0.05	<0.05	*IS*	*IS*	< 0.05	< 0.05
	DTX1 Free	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	DTX1 Total	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	DTX2 Free	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	DTX2 Total	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
3411	GYM	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	Homo-YTX	mg/kg WMB	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	<0.02	<0.02
	OA Free	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	OA Total	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	PnTx-G	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	PTX2	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	SPX1	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	Total DST	OA eq. mg/kg	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	YTX	mg/kg WMB	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
3411A	Brevetoxin 2	mg/kg WMB	<0.01*	*IS*	<0.01*	<0.01*	<0.01*	*IS*	<0.01*	<0.01*
	Brevetoxin 3	mg/kg WMB	0.01*	*IS*	0.09*	0.02*	0.02*	*IS*	0.03*	0.01*
	C1	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
3416	C2	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	<0.02	< 0.02
	СЗ	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	<0.02	<0.02
	C4	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	<0.02	<0.02
	dcGTX1	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02*	<0.02*	*IS*	*IS*	<0.02*	<0.02*
	dcGTX2	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	<0.02	<0.02
	dcGTX3	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	<0.02	< 0.02
	dcGTX4	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02*	<0.02*	*IS*	*IS*	<0.02*	<0.02*
	dcNEO	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	<0.02	<0.02

^{*}IS*- Insufficient Sample

^{*} NATA accreditation does not cover this result

Chemistry Test Results (Biota - Food)		Sample Description	Cormorant 1 Brain	Cormorant 1 Kidney	Cormorant 1	Cormorant 1 Lung	Cormorant 2 Brain	Cormorant 2 Kidney	Cormorant 2 Liver	Cormorant 2 Lung
		Sampled Date/ Time	07/07/25 0:00	07/07/25 0:00	07/07/25 0:00	07/07/25 0:00	07/07/25 0:00	07/07/25 0:00	07/07/25 0:00	07/07/25 0:00
Method ID	Analyte	Units	338829	338830	338831	338832	338833	338834	338835	338836
	dcSTX	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	doSTX	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.01*	<0.01*	*IS*	*IS*	<0.01*	<0.01*
	GTX1	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	GTX2	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	GTX3	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
3416	GTX4	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	GTX5	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	<0.02	<0.02
	GTX6	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	< 0.02	<0.02
	NEO	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.02	<0.02	*IS*	*IS*	<0.02	<0.02
	STX	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.01	<0.01	*IS*	*IS*	<0.01	<0.01
	Total PST	STX.2HCl eq. mg/kg	*IS*	*IS*	<0.10	<0.10	*IS*	*IS*	<0.10	<0.10

^{*}IS*- Insufficient Sample

^{*} NATA accreditation does not cover this result

Chemistry Test Results (Biota - Food)		Sample Description Sampled Date/ Time	Cormorant 3 Brain 07/07/25 0:00	Cormorant 3 Kidney 07/07/25 0:00	Cormorant 3 Liver 07/07/25 0:00	Cormorant 3 Lung 07/07/25 0:00	
Method ID	Analyte AZA1 AZA2 AZA3 Domoic Acid DTX1 Free DTX1 Total DTX2 Free DTX2 Total GYM Homo-YTX OA Free OA Total PnTx-G PTX2	Units	338837	338838	338839	338840	
	AZA1	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	AZA2	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	AZA3	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
3411	Domoic Acid	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	DTX1 Free	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	DTX1 Total	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	DTX2 Free	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	DTX2 Total	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	GYM	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	Homo-YTX	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	OA Free	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	OA Total	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	PnTx-G	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	PTX2	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	SPX1	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	Total DST	OA eq. mg/kg	*IS*	*IS*	*IS*	*IS*	
	YTX	mg/kg WMB	*IS*	*IS*	*IS*	*IS*	
	Brevetoxin 2	mg/kg WMB	<0.01*	*IS*	*IS*	<0.01*	
3411A	Brevetoxin 3	mg/kg WMB	<0.01*	*IS*	*IS*	0.01*	
	C1	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.01	
	C2	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.02	
	C3	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	< 0.02	
	C4	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	< 0.02	
3416	dcGTX1	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.02*	
	dcGTX2	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	< 0.02	
	dcGTX3	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	< 0.02	
	dcGTX4	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.02*	
	dcNEO	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	< 0.02	

^{*}IS*- Insufficient Sample

^{*} NATA accreditation does not cover this result

Chemistry Test Results (Biota - Food) Method ID Analyte		Sample Description Sampled Date/ Time	Cormorant 3 Brain 07/07/25 0:00	Cormorant 3 Kidney 07/07/25 0:00	Cormorant 3 Liver 07/07/25 0:00	Cormorant 3 Lung
		Units	338837	338838	338839	07/07/25 0:00 338840
	dcSTX	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.01
	doSTX	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.01*
	GTX1	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.01
	GTX2	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.01
	GTX3	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.01
3416	GTX4	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.01
	GTX5	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.02
	GTX6	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.02
	NEO	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.02
	STX	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.01
	Total PST	STX.2HCl eq. mg/kg	*IS*	*IS*	*IS*	<0.10

[&]quot;IS"- Insufficient Sample

^{*} NATA accreditation does not cover this result



PATH RESULTS: COMORANT MORTALITY, (Wi)

From

Date Fri 19/09/2025 4:00 PM

То



Owner: COMORANT MORTALITY KANGAROO ISLAND SHAOLS BEACH Animal/s: Wild Birds

Tested on

09/07/25

Referred on 19/09/25 16:30 **Referred on** 07/07/25 **by**:

DOB: N/A

Collected: 07/07/25 00:25 Subm.No:

Lab No.:

Samples tested as received All Te

bampies tested as rece

All Tests Complete

SUMMARY DIAGNOSIS Bird 2 Chronic hepatitis Renal trematodiasis

Birds 1-8 Alimentary ascardiasis

SUMMARY COMMENTS

The cause of chronic weight loss, neurological signs and death is not determined by laboratory tests. Avian influenza and Newcastle disease are excluded as causes of morbidity, neurological signs and death based on the negative PCR results.

Hepatitis contributed to morbidity for bird 2. Bacterial cultures are unremarkable. The renal trematode Renicola has been recorded in the kidneys of cormorants (P. Ladds 2009; Pathology of Australian native wildlife). This is an incidental finding.

The Cl. perfringens from bird 3 liver culture is a post mortem overgrowth.

Alimentary ascardiasis (Anisakis sp. or Contracaecum sp.) is a normal finding in marine birds.

ADDITIONAL COMMENTS 19/9/2025

Brevetoxin 3 was detected in the liver (0.09mg/kg), brain (0.01mg/kg) and lung (0.02mg/kg) of cormorant 1, brain (0.02mg/kg), liver (0.03mg/kg), lung (0.01mg/kg) of cormorant 2 and lung (0.01mg/kg) of cormorant 3. Based on the history of neurological signs in this flock of cormorants and the detection of brevetoxin 3, brevetoxin may have contributed morbidity for this flock. These findings should be interpreted in conjunction with any algal counts in the ocean waters near where they birds were found, around the time they showed neurological signs / were found dead.

Additional comments:

As little as 4.8 ng PbTx-3 eq./g, brevetoxin 3, and 3 ng PbTx-3 eq./g were detected by high performance liquid chromatography in the livers and 3 ng PbTx eq./kg in the brain from stranded brown pelicans exposed to Karenia brevis blooms along the central west coast of Florida, USA (Fauquier et al 2013). Fauquier and co-authors assessed that brevetoxin exposure was the probable cause of stranding in birds if they presented



Referred on 07/07/25 by

Owner:

COMORANT MORTALITY KANGAROO ISLAND SHAOLS BEACH Animal/s:

Wild Birds

DOB: N/A

Collected: 07/07/25 00:25 Subm.No:

Lab No.:

Samples tested as received

All Tests Complete

debilitated with detectable premortem or post mortem brevetoxin levels and had clinical neurological signs or stranded during the Karenia brevis bloom on the central west coast of Florida. Many of the affected birds on necropsy had chronic weight loss. There were no specific histopathological changes related to brevetoxicosis.

Fauquier, D.A., Flewelling, L.J., Maucher, J.M., Keller, M., Kinsel, M.J., Johnson, C.K., Henry, M., Gannon, J.G., Ramsdell, J.S. and Landsberg, J.H., 2013. Brevetoxicosis in seabirds naturally exposed to Karenia brevis blooms along the central west coast of Florida. Journal of wildlife diseases, 49(2), pp.246-260.

Specialist Veterinary Anatomic Pathologist

Validated by