

Inland Waters & Catchment Ecology

SOUTH
AUSTRALIAN
RESEARCH &
DEVELOPMENT
INSTITUTE
PIRSA

Lower Lakes Vegetation Condition Monitoring – 2012/2013



Kate Frahn, Susan Gehrig, Jason Nicol and Kelly Marsland

SARDI Publication No. F2009/000370-5
SARDI Research Report Series No. 707

SARDI Aquatic Sciences
PO Box 120 Henley Beach SA 5022

August 2013

Lower Lakes Vegetation Condition Monitoring – 2012/2013

Kate Frahn, Susan Gehrig, Jason Nicol and Kelly Marsland

**SARDI Publication No. F2009/000370-5
SARDI Research Report Series No. 707**

August 2013

This Publication may be cited as:

Frahn, K.A., Gehrig, S. L., Nicol, J.M., and Marsland, K.B. (2013). Lower Lakes vegetation condition monitoring – 2012/2013. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2009/000370-5. SARDI Research Report Series No. 707. 69pp.

South Australian Research and Development Institute

SARDI Aquatic Sciences

2 Hamra Avenue

West Beach SA 5024

Telephone: (08) 8207 2400

Facsimile: (08) 8207 5406

<http://www.sardi.sa.gov.au>

DISCLAIMER

SARDI Aquatic Sciences and the Department of Environment, Water and Natural Resources (DEWNR), do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaim all liability for any error, loss or other consequence which may arise from you relying on any information in this publication. The contents of the material do not purport to represent the position of the Commonwealth of Australia or the MDBA in any way and, as appropriate, are presented for the purpose of informing and stimulating discussion for improved management of Basin's natural resources. The report has been through the SARDI internal review process, and has been formally approved for release by the Research Chief, Aquatic Sciences. The SARDI Report Series is an Administrative Report Series which has not reviewed outside the department and is not considered peer-reviewed literature. Material presented in these Administrative Reports may later be published in formal peer-reviewed scientific literature.

© 2013 SARDI & DEWNR

This work is copyright. Apart from any use as permitted under the *Copyright Act* 1968 (Cth), no part may be reproduced by any process, electronic or otherwise, without the specific written permission of the copyright owner. Neither may information be stored electronically in any form whatsoever without such permission. This project was funded by The Living Murray initiative of the Murray- Darling Basin Commission, which has now transitioned to become the Murray Darling Basin Authority.

Printed in Adelaide: August 2013


SARDI Publication No. F2009/000370-5

SARDI Research Report Series No. 707

Authors: Kate Frahn, Susan Gehrig, Jason Nicol and Kelly Marsland

Reviewers: Chris Bice and Adrienne Frears

Approved by: Dr Qifeng Ye
Science Leader – Inland Waters & Catchment Ecology

Signed: 

Date: 28 August 2013

Distribution: MDBA, DEWNR, SAASC Library, University of Adelaide Library, Parliamentary Library, State Library and National Library

Circulation: Public Domain

Table of Contents

Table of Contents.....	i
List of Figures	i
List of Tables.....	iii
List of Appendices	iv
Acknowledgements	1
Executive Summary.....	2
1 Introduction.....	4
2 Methods.....	7
2.1 Study site, hydrology and salinity.....	7
2.2 Vegetation surveying protocol	12
2.3 Functional Groups.....	15
2.4 Data analysis.....	19
3 Results.....	20
3.1 Wetlands	20
3.2 Lakeshores.....	25
4 Discussion and management implications	41
4.1 Wetlands	41
4.2 Lakeshores.....	44
4.3 The Living Murray Target V3	47
4.4 Further studies	48
5 References	49
6 Appendices.....	56

List of Figures

Figure 1: Map of Lakes Alexandrina and Albert and Goolwa Channel showing the location of lakeshore and wetland vegetation monitoring sites (site numbers correspond to Table 1) and

major flow control structures (where sites are in close proximity they may not be visible on map).....9

Figure 2: Daily mean water levels in Goolwa Channel (Signal Point), Lake Alexandrina (Milang) and Lake Albert (Meningie) from August 2008 to April 2013 (except Goolwa Channel August 2008 to November 2012) (Department for Environment, Water and Natural Resources 2013b).10

Figure 3: Daily mean surface water electrical conductivity (EC) in Goolwa Channel (Signal Point), Lake Alexandrina (Milang) and Lake Albert (Meningie) from August 2008 to April 2013 (except Goolwa Channel August 2008 to November 2012) (Department for Environment, Water and Natural Resources 2013a).11

Figure 4: Vegetation surveying protocol for lakeshore sites: plan view showing placement of quadrats relative to the shoreline.14

Figure 5: Plant functional groups in relation to depth and duration of flooding.17

Figure 6: Proportion of plant functional groups for edge (A) and channel (B) elevations of Angas and Bremer River Mouths from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).27

Figure 7: Proportion of plant functional groups from highest (A: peg 5) to lowest elevation (E: peg 1) in Dunn's Lagoon from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).28

Figure 8: Proportion of plant functional groups from highest (A: peg 3) to lowest elevation (C: peg 1) in Goolwa Channel Drive from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).29

Figure 9: Proportion of plant functional groups for edge (A) and channel (B) elevations of Hunter's Creek from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).30

Figure 10: Proportion of plant functional groups across highest elevation (A: peg 4) to lowest elevation (E: peg 1) in Loveday Bay wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).31

Figure 11: Proportion of plant functional groups across highest (A: peg 1) to lowest elevation (D: peg 4) in Milang Wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).32

Figure 12: Proportion of plant functional groups across highest elevation (A: peg 4) to lowest elevation (D: peg 1) in Narrung wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).33

Figure 13: Proportion of plant functional groups across highest (A: peg 3) to lowest elevation (C: peg 1) in Point Sturt wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).34

Figure 14: Proportion of plant functional groups across highest (A: peg 4) to lowest elevation (D: peg 1) in Poltalloch wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).....	35
Figure 15: Proportion of plant functional groups across highest (A: peg 4) to lowest elevation (D: peg 1) in Teringie from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).	36
Figure 16: Proportion of plant functional groups across highest elevation (A: peg 4) to lowest elevation (D: peg 1) in Waltowa from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).....	37
Figure 17: Proportion of plant functional groups from highest elevation (+0.8 m AHD) to lowest elevation (-0.5 m AHD) in Lake Alexandrina from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).....	38
Figure 18: Proportion of plant functional groups from highest elevation (+0.8 m AHD) to lowest elevation (-0.5 m AHD) in Lake Albert from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).	39
Figure 19: Proportion of plant functional groups from highest elevation (+0.8 m AHD) to lowest elevation (-0.5 m AHD) in Goolwa Channel from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).....	40

List of Tables

Table 1: List of understorey vegetation site numbers (relative to map provided in Figure 1), site name, location, habitat type (wetland or lakeshore), number of survey sites and the year sites were established.	12
Table 2: Modified Braun-Blanquet (1932) scale estimating cover/abundance as per Heard and Channon (1997).	14
Table 3: Functional classification of plant species based on water regime preferences, modified from Brock and Casanova (1997) (*denotes exotic species).	16
Table 4: PERMANOVA results comparing the plant community through time and between elevations for each wetland.	20
Table 5: PERMANOVA results comparing the plant community through time and between elevations for Lake Alexandrina, Lake Albert and Goolwa Channel shorelines.	25
Table 6: Colour codes for vegetation functional groups.	27

List of Appendices

Appendix 1: GPS coordinates (UTM format, map datum WGS84) for lakeshore and wetland understorey vegetation monitoring sites.....	56
Appendix 2: Current nomenclature changes Centre for Australian National Biodiversity Research and Council of Heads of Australasian Herbaria (2013).	57
Appendix 3: Species list, functional classification (Gehrig and Nicol 2010b), life history strategy and conservation status (state conservation status from listings in Barker <i>et al.</i> (2005) and regional conservation status from listings in Lang and Kraehenuehl (2001) from all sites and survey dates (*denotes exotic taxon, **denotes proclaimed pest plant in South Australia, ***denotes weed of national significance # denotes listed as rare in South Australia).	58
Appendix 4: Taxa present (P) in each wetland from spring 2008 to autumn 2013 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).	62
Appendix 5: Taxa present (P) at lakeshore sites from spring 2008 to autumn 2013 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia)	67

Acknowledgements

The authors would like to thank Arron Strawbridge, Rod Ward, Kate Mason, Geoff Braddock, Robert Mrongovius, Margaret Mrongovius, Charles Andre, Jo Andre, John Rowe, Lorraine Rowe, Bev Stone, Alex Stone, Dennis Chandler, the Raukkan Natural Resources Management Team and Colleen Bakker for field assistance. Adrienne Frears, Rod Ward, Jason Tanner and Chris Bice for comments on early drafts of this report. Geoff and Libby Braddock, Dennis Chandler, Chris and Beth Cowan, the Raukkan Natural Resources Management Team, Charles and Jo Andre and David and Cathy Harvey for access to their properties. Geoff and Libby Braddock, Charles and Jo Andre, Alex and Bev Stone for lunches, tea, coffee and scones. Funding for condition monitoring has been provided by the Murray-Darling Basin Authority through The Living Murray initiative. This project has been managed by the Department of Environment, Water and Natural Resources, through the Lower Lakes, Coorong and Murray Mouth Icon Site staff.

Executive Summary

The Condition Monitoring Plan for the Coorong, Lower Lakes and Murray Mouth Icon Site (Maunsell Australia Pty Ltd. 2009) identified that a monitoring program was required that focused on measuring key environmental parameters that were considered to be indicators of river and floodplain health in both the short- and long-term. This report presents the findings of the first five years of a monitoring program established to evaluate The Living Murray Target V3 in the aforementioned plan: *maintain or improve aquatic and littoral vegetation in the Lower Lakes*.

Vegetation surveys were conducted at selected wetlands and lakeshore sites across Lakes Alexandrina and Albert, Goolwa Channel, the lower Finnis River, lower Currency Creek and the mouths of the Angas and Bremer Rivers. Sites established in spring 2008 and spring 2009 (Goolwa Channel monitoring sites) were re-surveyed. At each site, transects were established perpendicular to the shoreline and three, 1 x 3 m quadrats separated by 1 m were located at regular elevation intervals (defined by plant community) for wetlands or elevations (+0.8, +0.6, +0.4, +0.2, 0 and -0.5 m AHD) for lakeshores. The cover and abundance of each species present in quadrats were estimated using a modified Braun-Blanquet (1932) cover abundance score. Vegetation surveys were undertaken every spring (October 2008, 2009, November 2010, October 2011 and 2012) and autumn (March 2009, 2010, 2011, 2012 and 2013). The first two years of the monitoring program coincided with a period of record low water levels in the Lower Lakes. However, during this period, significant engineering interventions (construction of the Clayton Regulator and Narrung Bund and pumping of environmental water into Narrung Wetland) also influenced vegetation communities and were considered as part of the monitoring program. In August 2010, water levels in Lake Alexandrina rapidly rose to historical levels and in September 2010, the Clayton Regulator and Narrung Bund were breached, reconnecting these areas with Lake Alexandrina. High water levels, and subsequent connectivity, continued throughout the remainder of the condition monitoring program to date.

Over the five year period (spring 2008 to autumn 2013), a total of 145 taxa (including 79 exotics, three weeds of national significance, eight proclaimed pest plants in South Australia and one species listed as rare in South Australia) were recorded throughout the Lower Lakes. At wetland sites, 127 taxa were recorded (including 62 exotics) and at lakeshore sites 106 taxa were recorded (63 exotics). From 2008 to 2010, disconnection and subsequent desiccation of wetlands generally resulted in the loss of submergent taxa (excepting wetlands that received rainfall runoff or were filled by pumping i.e. Narrung) and colonisation by terrestrial species. In 2009-10, more terrestrial species had colonised lower elevations than previously recorded in 2008-09. When water levels were reinstated, aquatic species generally recolonised the inundated areas and increased in cover for the remainder of the survey period, albeit not to the extent recorded in

2004 (Holt *et al.* 2005) and 2005 (Nicol *et al.* 2006). Waltowa wetland remained dominated by salt tolerant and terrestrial taxa despite active management of water levels from May 2011 to April 2013.

Similar to wetlands, low water levels resulted in colonisation of terrestrial species around the shorelines of Lakes Alexandrina and Albert (lower elevations were colonised in 2009-10 compared with 2008-09). In contrast, by March 2010, an amphibious, emergent and submergent plant community developed in areas inundated by the Clayton Regulator (which in 2008-09 were dominated by terrestrial taxa and bare soil). Water level management in Lake Albert did not result in colonisation of aquatic species or reconnect fringing vegetation. When water levels were reinstated, terrestrial taxa were extirpated in Lakes Alexandrina and Albert. Furthermore, there was an increase in the abundance of emergent, amphibious and submergent species in Lake Alexandrina and emergent species in Lake Albert. Breaching of the Clayton Regulator decreased salinity in Goolwa Channel, which resulted in a further change to the plant community. The submergent species, *Potamogeton pectinatus*, dominated large areas of Goolwa Channel from autumn 2010 to spring 2010 but, was replaced by *Myriophyllum salsugineum* by autumn 2011 and a more diverse submergent plant community that also included *Vallisneria australis*, *Ceratophyllum demersum*, *Potamogeton crispus* and *Myriophyllum caput-medusae* was present in March 2013. The emergent plant community in Goolwa Channel also changed significantly over the same period with *Schoenoplectus validus* abundance increasing between spring 2010 and autumn 2011. In 2011/12, the aquatic and emergent plant community recovered further, although the changes were less dramatic than observed in the previous 12 months and in 2012/13 there was little change in the emergent plant community.

Results showed that the plant community is resilient and recovering but the submergent plant community is different (cover and diversity) from the community present before water levels were drawn down. Large areas that were historically dominated by diverse submergent plant communities (e.g. Clayton Bay, Dunns Lagoon, Teringie and Narrung) are yet to fully recover. *Myriophyllum caput-medusae* was recorded for the first time since 2005; however, *Lepilaena cylindrocarpa* and *Ranunculus trichophyllus* have not been recorded.

Comparison between monitoring results and the River Murray Wetlands baseline surveys undertaken in 2004 and 2005 show that target V3 has still not been met for understorey vegetation, except in Goolwa Channel, the lower Finniss River and lower Currency Creek. However, after water levels increased there was a significant and sustained increase in the abundance, extent and diversity of aquatic plant communities. In addition, all of the aquatic species that have recruited are capable of rapidly colonising large areas by asexual reproduction providing the current hydrological and salinity regime is maintained. Therefore, TLM target V3 may be attained for the Lower Lakes icon site in the near future.

1 Introduction

The Coorong, Lower Lakes and Murray Mouth region has been listed as one of six icon sites under the Murray-Darling Basin Authority's "The Living Murray" (TLM) program and has been identified as an indicator site under the "Basin Plan". The Condition Monitoring Plan for the Coorong, Lower Lakes and Murray Mouth Icon Site outlined a series of 17 condition targets for the Icon Site (Maunsell Australia Pty Ltd. 2009). This report presents the findings from the first five years of the understorey component of a condition monitoring program designed to evaluate target V3: *maintain or improve aquatic and littoral vegetation in the Lower Lakes* (Marsland and Nicol 2009; Gehrig *et al.* 2010; 2011b; 2012).

Scientifically defensible and statistically robust monitoring programs need to be established to assist in meeting the ecological targets in the Coorong, Lower Lakes and Murray Mouth Icon Site Environmental Management Plan and Ramsar Management Plan. Marsland and Nicol (2006) identified that existing monitoring programs (in 2006) would not adequately assess target V3; therefore, a monitoring program that expanded and built upon existing monitoring programs (South Australian Murray-Darling Basin Natural Resources Management (SAMDBNRM) Board community wetland monitoring) was established in 2008 (Marsland and Nicol 2009). The understorey vegetation monitoring program described in this report uses the same methods and sites as the SAMDBNRM Board community wetland monitoring program but includes additional sites in lakeshore habitats (in Lakes Alexandrina and Albert), the lower reaches of the Finnis River, Currency Creek and Goolwa Channel and wetlands that were not part of the original community wetland monitoring program (Marsland and Nicol 2009). In 2009, eight extra sites in Goolwa Channel were added to assess the impact of the Goolwa Channel Water Level Management Project, and data from this project was included in the TLM Condition Monitoring Program (Gehrig and Nicol 2010a; Gehrig *et al.* 2010; 2011a; 2012).

The Condition Monitoring Plan for the Icon Site proposed 'indicators for monitoring' that comprised of individual taxa and discrete communities: *Melaleuca halmaturorum*, *Myriophyllum* spp., *Gabnia filum*, *Schoenoplectus* spp., *Typha domingensis*, *Phragmites australis* and samphire communities (Maunsell Australia Pty Ltd. 2009). However, further discussions concluded that the entire understorey vegetation assemblage would be monitored with a separate technique used for the dominant tree species *Melaleuca halmaturorum*. Hence, the monitoring program consists of two complementary components: the first component involves the monitoring of aquatic and littoral understorey vegetation in spring (high lake levels) and autumn (low lake levels) to determine the current condition, seasonal changes and medium- to long-term changes in floristic composition, and the second component monitors the mid- to long-term population dynamics of *Melaleuca halmaturorum*. The *Melaleuca halmaturorum* component of the monitoring program was only

undertaken in 2008/09; hence stand condition was not monitored in 2012-13. Information regarding *Melaleuca balmaturorum* stand condition is presented in Marsland and Nicol (2009).

From 1996 to 2010, the Murray-Darling Basin was subjected to the most severe drought in recorded history (Van Dijk *et al.* 2013). Below average stream flows coupled with upstream extraction and river regulation resulted in reduced inflows into South Australia (Van Dijk *et al.* 2013), which between January 2007 and August 2010 were insufficient to maintain pool level downstream of Lock and Weir number 1. Subsequently water levels in Lakes Alexandrina and Albert dropped to unprecedented lows (<-0.75 m AHD), fringing wetlands became disconnected and extensive areas of acid sulfate soils were exposed; particularly in Lake Albert and the lower reaches of the Finniss River and Currency Creek (Merry *et al.* 2003; Fitzpatrick *et al.* 2009a; Fitzpatrick *et al.* 2009b).

Prior to 2007, fringing wetlands in the Lower Lakes region contained diverse communities of emergent, amphibious and submergent taxa (Renfrey *et al.* 1989; Holt *et al.* 2005; Nicol *et al.* 2006). By spring 2008, submergent taxa had been extirpated (except for a small number of *Ruppia tuberosa* plants in Hunters Creek, in Lake Alexandrina near Raukkun and Loveday Bay Wetland and *Lamprothamnium macropogon* in Loveday Bay Wetland), amphibious taxa had declined in abundance and diversity, stands of emergent taxa were disconnected from remaining water and fringing habitats were dominated by terrestrial taxa and bare soil (Marsland and Nicol 2009). Furthermore, submergent taxa had not colonised the remaining open water areas (Marsland and Nicol 2009).

The loss of submergent vegetation, decline in abundance and diversity of amphibious taxa and disconnection of fringing emergent macrophytes had serious implications for ecosystem dynamics of the Lower Lakes. Aquatic vegetation is a critical ecosystem component in the Lower Lakes; plants are major primary producers (e.g. dos Santos and Esteves 2002; Camargo *et al.* 2006; Noges *et al.* 2010), improve water quality (e.g. Webster *et al.* 2001; James *et al.* 2004) provide habitat for invertebrates (e.g. Declerck *et al.* 2005; Cronin *et al.* 2006; Pinto *et al.* 2006; Papas 2007), birds (e.g. Brandle *et al.* 2002; Phillips and Muller 2006) and threatened fish (Wedderburn *et al.* 2007; Bice *et al.* 2008) and stabilise shorelines (Abernethy and Rutherford 1998; PIRSA Spatial Information Services 2009).

To mitigate acid sulfate soils three regulators were constructed in the Lower Lakes: the Narrung Bund, the Clayton Regulator and the Currency Creek Regulator (Figure 1). However, only the impacts of the Narrung Bund and Clayton Regulator will be discussed in this report due to the Currency Creek Regulator spillway remaining inundated after the Clayton regulator was constructed. The regulators disconnected Goolwa Channel and Lake Albert from Lake Alexandrina, which enabled water levels within each site to be managed independently. An

additional hydrological intervention was undertaken at Narrung Wetland, with 250 ML of environmental water from Lake Alexandrina being pumped into the wetland in October 2009 to provide suitable conditions for the growth of submergent taxa (particularly *Ruppia tuberosa*).

In August 2010, flows into South Australia increased, water levels in Lake Alexandrina were reinstated to historical levels ($\sim +0.75$ m AHD) and there was significant flow through the Barrages for the first time since spring 2005 (although there was a small release in 2006-07 to operate fishways). Furthermore, the Clayton Regulator and Narrung Bund were breached in September 2010, and Lake Alexandrina was reconnected with Goolwa Channel and Lake Albert. The impacts of the regulators, pumping and unregulated River Murray flows on salinity and water levels are outlined in section 2.1

The recent period of low flow, regulator construction, pumping, unregulated River Murray flows and regulator breaching have resulted in large changes to the hydrological and salinity regime of the Lower Lakes since 2007. Salinity (e.g. Hart *et al.* 2003; James *et al.* 2003; Nielsen *et al.* 2003; Nielsen and Brock 2009) and water regime (determined by lake levels) (e.g. Brownlow 1997; Blanch *et al.* 1999; Brock *et al.* 1999; Blanch *et al.* 2000; Nicol *et al.* 2003) are two of the primary drivers of plant community composition in freshwater ecosystems. Historically, the various components of the system were connected with relatively stable water levels ranging from +0.4 to +0.8 m AHD and surface water electrical conductivity lower than $2,000 \mu\text{S}\cdot\text{cm}^{-1}$ (Kingsford *et al.* 2009; 2011). Between 2007 and August 2010, surface water salinity, water regime and connectivity of the study area varied dramatically from historical patterns; however, since September 2010 the aforementioned factors have reflected historical patterns, except in Lake Albert where salinities have remained elevated.

The monitoring undertaken in 2012-13 builds on data collected between 2008 and 2012 and provides information regarding the change in plant communities since spring 2008. The survey period includes a period of record low water levels in Lake Alexandrina, several engineering interventions, an unregulated River Murray flow, two in channel flow pulses and entitlement flows. Therefore, this monitoring program collected information regarding the change in wetland plant communities in response to drawdown, desiccation and increased water levels due to regulated inundation, and natural flooding, and provides an insight into recovery of the system under hydrological restoration. The aims of this project were to:

- Continue the statistically robust, quantitative understorey aquatic and littoral vegetation monitoring program in the Lower Lakes to assess TLM target V3.
- Monitor the early stages of recovery of the aquatic plant community after hydrological restoration following extended drought, drawdown, fragmentation and desiccation of aquatic habitats.

2 Methods

2.1 Study site, hydrology and salinity

Vegetation surveys were undertaken in Goolwa Channel, the lower Finnis River, Lower Currency Creek (herein referred to collectively as the Goolwa Channel), Lake Alexandrina and Lake Albert (Figure 1). Since early 2008, a range of interventions have been undertaken in the Lower Lakes to regulate water levels and mitigate acid sulfate soils; primarily being the construction of the Narrung Bund and Clayton Regulator (Figure 1). Construction of the Narrung Bund was completed in early 2008 and disconnected Lake Albert from Lake Alexandrina (Figure 1). Water was then pumped from Lake Alexandrina into Lake Albert to maintain water levels above -0.5 m AHD. Construction of the Clayton Regulator was completed in August 2009 and impounded water from the Finnis River and Currency and Tookayerta Creeks (Figure 1). In addition, water was pumped into Goolwa Channel (Figure 2) from Lake Alexandrina to raise water levels to +0.7 m AHD in spring 2009.

Construction of the Narrung Bund and Clayton Regulator enabled independent management of Lake Alexandrina, Lake Albert and Goolwa Channel, which is reflected in water levels (Figure 2) and surface water salinity (measured as electrical conductivity (EC, $\mu\text{S.cm}^{-1}$) (Figure 3). Water levels in Lake Alexandrina were dependent on River Murray inflows and, to a lesser extent, pumping into Lake Albert and Goolwa Channel. Due to low flows into South Australia, the water level in Lake Alexandrina remained below sea level for much of this previous survey period (Figure 2). River Murray flows into Lake Alexandrina increased in April 2010, and the lake was restored to historical water levels in August 2010 and fluctuated between +0.4 and +0.8 m AHD for the remainder of the survey period (Figure 2). Whilst River Murray inflows were below average, surface water EC in Lake Alexandrina remained relatively constant and ranged from 4,000 to 7,000 $\mu\text{S.cm}^{-1}$ (Figure 3). When inflows increased, EC decreased and by December 2010 had fallen to around 500 $\mu\text{S.cm}^{-1}$ for the remainder of the survey period (Figure 3).

From August 2008 to August 2010, water levels in Lake Albert were dependent on pumping from Lake Alexandrina, local rainfall and evaporation. Between August 2008 and March 2009, water level decreased from -0.1 to -0.55 m AHD then increased to approximately -0.1 m AHD by September 2009 (Figure 2) as a result of pumping from Lake Alexandrina. Pumping ceased in September 2009 and water levels decreased to -0.7 m AHD in January 2010 (Figure 2). Pumping recommenced between April and June 2010 and water levels increased to -0.4 m AHD (Figure 2). In September 2010, the Narrung Bund was breached and Lake Albert was reconnected with Lake Alexandrina. Water level increased rapidly to +0.8 m AHD, reflecting water level in Lake Alexandrina and ranging between +0.4 and +0.8 m AHD for the remainder of the study period

(Figure 2). Surface water EC increased from 5,000 to 12,000 $\mu\text{S.cm}^{-1}$ from August 2008 to March 2009, then (as a result of pumping from Lake Alexandrina), decreased to around 9,500 $\mu\text{S.cm}^{-1}$ by May 2009, remaining relatively constant until October 2009 (Figure 3). When pumping ceased, surface water EC increased and exceeded 20,000 $\mu\text{S.cm}^{-1}$ by February 2010 (Figure 3). When pumping recommenced EC decreased to approximately 14,000 $\mu\text{S.cm}^{-1}$ until September 2010 when the bund was breached (Figure 3). After the bund was breached, EC decreased rapidly to 8,000 $\mu\text{S.cm}^{-1}$ and then slowly decreased to approximately 3,000 $\mu\text{S.cm}^{-1}$ by autumn 2013 (Figure 3).

Water levels in Goolwa Channel from August 2008 to August 2009 reflected those in Lake Alexandrina (Figure 2). From August 2009 to August 2010, following completion of the Clayton Regulator, water level was dependent upon pumping from Lake Alexandrina, inflows from the Finnis River and Currency and Tookayerta Creeks, local rainfall and evaporation and reached +0.75 m AHD in spring 2009 (Figure 2). Pumping ceased in November 2009 and water levels decreased to -0.1 m AHD in April/May 2010 (Figure 2). Water levels increased to +0.2 m AHD in response to tributary inflows in July 2010, and the regulator was breached in September 2010, which resulted in water levels rising to +0.8 m AHD (Figure 2). For the remainder of the survey period water levels were dependent on Lake Alexandrina water levels, but tended to fluctuate a little more (compared to Lake Alexandrina), ranging from +0.75 to +0.3 m AHD (no data available from November 2012 onwards) (Figure 2). Surface water EC at the beginning of the study period was approximately 21,000 $\mu\text{S.cm}^{-1}$, which decreased to around 14,000 $\mu\text{S.cm}^{-1}$ in September 2008, and then increased over spring and summer reaching a maximum of 33,000 $\mu\text{S.cm}^{-1}$ in February/March 2009 (Figure 3). The elevated EC was due to a combination of low River Murray inflows and seawater leaking through Goolwa Barrage into Goolwa Channel. From March 2009 to August 2009, EC decreased to around 20,000 $\mu\text{S.cm}^{-1}$ (Figure 3) due to higher water levels in Lake Alexandrina and engineering works that reduced seawater leakage through Goolwa Barrage. Construction of the Clayton Regulator was completed and pumping from Lake Alexandrina commenced in August 2009, which reduced EC in Goolwa Channel to 10,000 $\mu\text{S.cm}^{-1}$ until pumping ceased in November 2009 (Figure 3). Surface water EC increased to approximately 20,000 $\mu\text{S.cm}^{-1}$ by April 2010 and fluctuated between 20,000 and 22,000 $\mu\text{S.cm}^{-1}$ until the regulator was breached in September 2010 (Figure 3). Post August 2010, EC in the Goolwa Channel decreased rapidly to around 500 $\mu\text{S.cm}^{-1}$ for the remainder of the survey period, except for several salinity spikes (maximum approximately 47,000 $\mu\text{S.cm}^{-1}$) between May 2011 and October 2012 as a result of very high tides pushing seawater through open barrage gates (no data available from November 2012 onwards) (Figure 3).




Figure 1: Map of Lakes Alexandrina and Albert and Goolwa Channel showing the location of lakeshore and wetland vegetation monitoring sites (site numbers correspond to Table 1) and major flow control structures (where sites are in close proximity they may not be visible on map).




Figure 2: Daily mean water levels in Goolwa Channel (Signal Point), Lake Alexandrina (Milang) and Lake Albert (Meningie) from August 2008 to April 2013 (except Goolwa Channel August 2008 to November 2012) (Department for Environment, Water and Natural Resources 2013b).




Figure 3: Daily mean surface water electrical conductivity (EC) in Goolwa Channel (Signal Point), Lake Alexandrina (Milang) and Lake Albert (Meningie) from August 2008 to April 2013 (except Goolwa Channel August 2008 to November 2012) (Department for Environment, Water and Natural Resources 2013a).

2.2 Vegetation surveying protocol

Monitoring of understorey vegetation was conducted at 11 wetland and 25 lakeshore sites in every spring and autumn from October 2008 to March 2013 (for sites established in 2008 or earlier) and every spring and autumn from October 2009 to March 2013 for sites established in 2009 (Table 1). Sites were grouped on the basis of habitat (lakeshore or wetland) and location (Lake Alexandrina, Lake Albert or Goolwa Channel). GPS coordinates for each site are listed in (Appendix 1).

Table 1: List of understorey vegetation site numbers (relative to map provided in Figure 1), site name, location, habitat type (wetland or lakeshore), number of survey sites and the year sites were established.

Site #	Site Name	Location	Habitat	No. Survey Sites	Year Established
1	Bremer Mouth Lakeshore	Lake Alexandrina	lakeshore	1	2008
2	Brown Beach 1	Lake Albert	lakeshore	1	2008
3	Brown Beach 2	Lake Albert	lakeshore	1	2008
4	Clayton Bay	Goolwa Channel	lakeshore	1	2009
5	Currency Creek 3	Goolwa Channel	lakeshore	1	2008
6	Currency Creek 4	Goolwa Channel	lakeshore	1	2008
7	Goolwa North	Goolwa Channel	lakeshore	1	2009
8	Goolwa South	Goolwa Channel	lakeshore	1	2009
9	Hindmarsh Island Bridge 01	Goolwa Channel	lakeshore	1	2009
10	Hindmarsh Island Bridge 02	Goolwa Channel	lakeshore	1	2009
11	Lake Reserve Rd	Lake Alexandrina	lakeshore	1	2008
12	Loveday Bay	Lake Alexandrina	lakeshore	1	2009
13	Loveday Bay Lakeshore	Lake Alexandrina	lakeshore	1	2009
14	Lower Finniss 02	Goolwa Channel	lakeshore	1	2009
15	Milang (existing SAMDBNRM Board community monitoring site)	Lake Alexandrina	wetland	1	pre-2008
16	Milang Lakeshore	Lake Alexandrina	lakeshore	1	2009
17	Pt Sturt Lakeshore	Lake Alexandrina	lakeshore	1	2008
18	Pt Sturt Water Reserve	Lake Alexandrina	lakeshore	1	2008
19	Teringie Lakeshore	Lake Alexandrina	lakeshore	1	2008
20	Upstream of Clayton Regulator	Lake Alexandrina	lakeshore	1	2009
21	Wally's Landing	Goolwa Channel	lakeshore	1	2009
22	Warrengie 1	Lake Albert	lakeshore	1	2009
23	Lower Finniss 03	Goolwa Channel	lakeshore	1	2009
24	Narrung Lakeshore	Lake Alexandrina	lakeshore	1	2008
25	Nurra Nurra	Lake Albert	lakeshore	1	2008
26	Warrengie 2	Lake Albert	lakeshore	1	2009
27	Angas Mouth	Lake Alexandrina	wetland	1	2008
28	Bremer Mouth	Lake Alexandrina	wetland	1	2008

Site #	Site Name	Location	Habitat	No. Survey Sites	Year Established
29	Dunns Lagoon	Lake Alexandrina	wetland	4	2008
30	Goolwa Channel Drive	Lake Alexandrina	wetland	3	2008
31	Hunters Creek	Lake Alexandrina	wetland	5	2008
32	Poltalloch	Lake Alexandrina	wetland	2	2008
33	Pt Sturt	Lake Alexandrina	wetland	2	2008
34	Teringie (existing SAMDBNRM Board community monitoring site)	Lake Alexandrina	wetland	4	pre-2008
35	Waltowa (existing SAMDBNRM Board community monitoring site)	Lake Albert	wetland	2	pre-2008
36	Narrung (existing SAMDBNRM Board community monitoring site)	Lake Alexandrina	wetland	3	pre-2008

2.2.1 Wetlands

At each survey site (Table 1), a transect running perpendicular to the shoreline and three, 1 x 3 m quadrats, separated by 1 m, were established (Figure 4) at regular elevation intervals that represented the dominant plant communities (A. Frears pers. comm.). In wetlands with an established monitoring program (Milang, Waltowa, Teringie and Narrung), existing sites were re-surveyed. For the remaining wetlands (Dunns Lagoon, Pt Sturt, Hunters Creek, Goolwa Channel Drive, Bremer River Mouth, Angas River Mouth and Loveday Bay), a transect was established and quadrats placed in each plant community present during the spring 2008 survey. A minimum of one additional transect (but usually two or more in each wetland, except the Angas and Bremer River mouths) was established, and quadrats were placed at the same elevations (determined using a laser level) as on the first transect. At sites where the elevation gradient was steep (e.g. Angas and Bremer River Mouth, Hunter's Creek) only edge and channel quadrats were surveyed. Cover and abundance of each species present in the quadrat were estimated using the method outlined in Heard and Channon (1997), except that N and T were replaced by 0.1 and 0.5 to enable statistical analyses (Table 2).

Table 2: Modified Braun-Blanquet (1932) scale estimating cover/abundance as per Heard and Channon (1997).

Score	Modified Score	Description
N	0.1	Not many, 1-10 individuals
T	0.5	Sparsely or very sparsely present; cover very small (less than 5%)
1	1	Plentiful but of small cover (less than 5%)
2	2	Any number of individuals covering 5-25% of the area
3	3	Any number of individuals covering 25-50% of the area
4	4	Any number of individuals covering 50-75% of the area
5	5	Covering more than 75% of the area

2.2.2 Lakeshores

At each site, a transect running perpendicular to the shoreline was established and three, 1 x 3 m quadrats, separated by 1 m, were established at elevation intervals of +0.8, +0.6, +0.4, +0.2, 0 and -0.5 m AHD (Figure 4) (*sensu* Marsland and Nicol 2009; Gehrig and Nicol 2010a; Gehrig *et al.* 2010; Nicol and Marsland 2010).

**Figure 4:** Vegetation surveying protocol for lakeshore sites: plan view showing placement of quadrats relative to the shoreline.

2.2.3 Plant identification and nomenclature

Plants were identified using keys in Cunningham *et al.* (1981), Jessop and Toelken (1986), Dashorst and Jessop (1998), Romanowski (1998), Sainty and Jacobs (1981; 2003), Prescott (1988) and Jessop *et al.* (2006). In some cases due to immature individuals or lack of floral structures plants were identified to genus only. Nomenclature follows the Centre for Australian National Biodiversity Research and Council of Heads of Australasian Herbaria (2013) and changes to nomenclature (relative to previous reports) are found in Appendix 2.

2.3 Functional Groups

Due to the large number of species present, species were classified into functional groups (based on water regime preferences) outlined in Table 3 (also see Appendix 3). The position each group occupies in relation to flooding depth and duration is outlined in Figure 4. The functional classification was based on the classification framework devised by Brock and Casanova (1997), which was based on species from wetlands in the New England Tablelands region of New South Wales and modified by Gehrig and Nicol (2010b) to suit the plant communities of the Lower Lakes.

The use of a functional group approach to assess change through time and potential impacts of management strategies has several advantages compared to a species or community based approach:

- species with similar water regime preferences are grouped together, which simplifies systems with high species richness (especially where there are large numbers of species with similar water regime preferences),
- predictions about the response of the plant community are made based on processes and does not require prior biological knowledge of the system,
- it is transferrable between systems,
- robust and testable models that predict the response of a system to an intervention or natural event can be constructed, which can in turn be used as hypotheses for monitoring programs.

However there are limitations to the approach, which include:

- loss of information on species or communities (especially if there are species or communities of conservation significance or there is a pest plant problem),
- uncertainty regarding which species should be classified into which functional group,

- important factors (e.g. salinity) are often not taken into consideration (additional factors can be included; however, this can often complicate the functional classification and in systems where there is low species richness the number of groups may be greater than the number of species).

In this report changes through time and between locations and elevations, and TLM targets will be assessed and discussed using both species and functional approaches.

Table 3: Functional classification of plant species based on water regime preferences, modified from Brock and Casanova (1997) (*denotes exotic species).

Functional Group	Water Regime Preference	Examples
Terrestrial dry	Will not tolerate inundation and tolerates low soil moisture for extended periods.	<i>Medicago</i> spp.* <i>Brassica rapa</i> * <i>Bromus</i> spp.*
Terrestrial damp	Will tolerate inundation for short periods (<2 weeks) but require high soil moisture throughout their life cycle.	<i>Centaurea calcitrapa</i> * <i>Chenopodium album</i> * <i>Fumaria bastardii</i> *
Floodplain	Temporary inundation, plants germinate on newly exposed soil after flooding but not in response to rainfall.	<i>Lachnagrostis filiformis</i>
Amphibious fluctuation tolerator-emergent	Fluctuating water levels, plants do not respond morphologically to flooding and drying and will tolerate short-term complete submergence (<2 weeks).	<i>Cyperus gymnocaulos</i> <i>Juncus kraussii</i> <i>Schoenoplectus pungens</i>
Amphibious fluctuation tolerator-woody	Fluctuating water levels, plants do not respond morphologically to flooding and drying and are large perennial woody species.	<i>Melaleuca halmaturorum</i> <i>Duma florulenta</i>
Amphibious fluctuation tolerator-low growing	Fluctuating water levels, plants do not respond morphologically to flooding and drying and are generally small herbaceous species.	<i>Isolepis producta</i> <i>Isolepis platycarpa</i>
Amphibious fluctuation responder-plastic	Fluctuating water levels, plants respond morphologically to flooding and drying (e.g. increasing above to below ground biomass ratios when flooded).	<i>Persicaria lapathifolia</i> <i>Ludwigia peploides</i> <i>Cotula coronopifolia</i> <i>Hydrocotyle verticillata</i>
Floating	Static or fluctuating water levels, plants respond to fluctuating water levels by having some or all organs floating on the water surface. Most species require permanent water to survive.	<i>Azolla</i> spp. <i>Lemna</i> spp.
Submergent r-selected	Temporary wetlands that hold water for longer than 4 months.	<i>Ruppia tuberosa</i> <i>Ruppia polycarpa</i>
Emergent	Static shallow water <1 m or permanently saturated soil.	<i>Typha</i> spp. <i>Phragmites australis</i> <i>Schoenoplectus validus</i>
Submergent k-selected	Permanent water.	<i>Myriophyllum salsugineum</i> <i>Vallisneria australis</i> <i>Ruppia megacarpa</i> <i>Potamogeton pectinatus</i>




Figure 5: Plant functional groups in relation to depth and duration of flooding.

The “terrestrial dry” functional group is intolerant of flooding and taxa will persist in environments with low soil moisture (Table 3) (Brock and Casanova 1997). Taxa from this functional group often invade wetlands that have been drawn down for an extended period or floodplains where there has been a lack of flooding, but are generally restricted to highlands that never flood (Brock and Casanova 1997).

Taxa in the “terrestrial damp” group will tolerate inundation for short periods and require high soil moisture to complete their life cycle (Table 3) (Brock and Casanova 1997). Taxa from this functional group are often winter annuals, perennial species that grow around the edges of permanent water bodies where there is high soil moisture or species that colonise wetlands shortly after they are drawn down and riparian zones and floodplains shortly after flood waters recede (Brock and Casanova 1997).

Taxa in the “floodplain” functional group exhibit most of the traits of terrestrial species; they are generally intolerant of long-term inundation but are restricted to areas that flood periodically (they are absent from the highlands) because they only germinate after flood waters recede or wetlands are drawn down, not in response to rainfall (Table 3) (Nicol 2004). Taxa from this functional group colonise floodplains and riparian zones after flood waters have receded and when wetlands are drawn down (Nicol 2004). Floodplain species often have flexible life history strategies; they grow whilst soil moisture is high and flower and set seed (after which most species die) in response to low soil moisture (Nicol 2004).

The “amphibious fluctuation tolerator-emergent” group consists mainly of emergent sedges and rushes that prefer high soil moisture or shallow water but require their photosynthetic parts to be emergent, although many will often tolerate short-term submergence (Table 3) (Brock and Casanova 1997). Taxa from this group are often found on the edges of permanent water bodies, in seasonal and temporary wetlands, in riparian zones and areas that frequently wet and dry.

Taxa in the “amphibious fluctuation tolerator-woody” group have similar water regime preferences to the amphibious fluctuation tolerator-emergent group (Figure 5) and consist of woody perennial species (Table 3) (Brock and Casanova 1997). Plants generally require high soil moisture in the root zone but there are several species that are tolerant of desiccation for extended periods (Roberts and Marston 2000). Taxa in this functional group are generally found on the edges of permanent water bodies, in seasonal and temporary wetlands, in riparian zones and areas that frequently wet and dry.

The “amphibious fluctuation tolerator-low growing” group have similar water regime preferences to the amphibious fluctuation tolerator-emergent and amphibious fluctuation tolerator-woody groups (Figure 5); however, some species can grow totally submerged except during flowering (when there is a requirement for a dry phase) (Table 3) (Brock and Casanova 1997). Taxa in this functional group are generally found on the edges of permanent water bodies, in seasonal and temporary wetlands, in riparian zones and areas that frequently wet and dry but taxa are usually less desiccation tolerant than species in the other amphibious tolerator groups (Table 3).

The “amphibious fluctuation responder-plastic” group occupies a similar zone to the amphibious fluctuation tolerator-low growing group; except that they have a physical response to water level changes such as rapid shoot elongation or a change in leaf type (Brock and Casanova 1997). They can persist on damp and drying ground because of their morphological flexibility but can flower even if the site does not dry out. They occupy a slightly deeper/wet for longer area than the amphibious fluctuation tolerator-low growing group (Figure 5).

Taxa in the “floating” functional group float on the top of the water (often unattached to the sediment) with the majority of species requiring the presence of free water of some depth year round; although, some species can survive and complete their life cycle stranded on mud (Table 3) (Brock and Casanova 1997). Taxa in this group are usually found in permanent waterbodies, often forming large floating mats upstream of barriers (e.g. weirs), in lentic water bodies and slackwaters.

“Submergent r-selected” taxa colonise recently flooded areas (Table 3) and show many of the attributes of Grime’s (1979) r-selected (ruderal) species, which are adapted to periodic disturbances. Many require drying to stimulate germination; they frequently complete their life cycle quickly and die off naturally. They persist via a dormant, long-lived bank of seeds, spores or asexual propagules (e.g. *Ruppia tuberosa* and *Ruppia polycarpa* turions in the sediment) (Brock 1982). They prefer habitats that are annually flooded to a

depth of more than 10cm but can persist as dormant propagules for a number of years (temporary or ephemeral wetlands).

The “emergent” group consists of taxa that require permanent shallow water or a permanently saturated root zone, but have emergent leaves or stems (Table 3). They are often found on the edges of permanent waterbodies and in permanent water up to 2 m deep (depending on species) or in areas where there are shallow water tables (Roberts and Marston 2000).

“Submergent k-selected” taxa require permanent water greater than 10 cm deep for more than a year to either germinate or reach sufficient biomass to start reproducing (Table 3) (Roberts and Marston 2000). Taxa in this group show many of the attributes of Grime’s (1979) k-selected (competitor) taxa that are adapted to stable environments and are only found in permanent water bodies. The depth of colonisation of submergent k-selected taxa is dependent on photosynthetic efficiency and water clarity (*sensu* Spence 1982).

2.4 Data analysis

2.4.1 Wetlands

Changes in floristic composition through time and between elevations were analysed for each wetland (except Angas and Bremer Mouths, where data were combined and treated as one wetland) using two-factor PERMANOVA (Anderson 2001; Anderson and Ter Braak 2003) using the package PRIMER version 6.1.12 (Clarke and Gorley 2006) and Indicator Species Analysis (Dufrene and Legendre 1997) using the package PCOrd version 5.12 (McCune and Mefford 2006).

2.4.2 Lakeshores

Lakeshore sites were analysed separately to the wetlands. Changes in floristic composition through time and between elevations for each location (Goolwa Channel, Lake Alexandrina and Lake Albert) were analysed independently using two-factor PERMANOVA (Anderson 2001; Anderson and Ter Braak 2003), using the package PRIMER version 6.1.12. (Clarke and Gorley 2006) and Indicator Species Analysis (Dufrene and Legendre 1997) using the package PCOrd version 5.12 (McCune and Mefford 2006). Bray-Curtis (1957) similarities were used to calculate the similarity matrices for all multivariate analyses. $\alpha=0.05$ was used for all statistical analyses.

3 Results

3.1 Wetlands

A total of 127 taxa (62 exotics including two weeds of national significance, seven proclaimed pest plants in South Australia and one species listed as rare in South Australia) were recorded for all wetland sites between spring 2008 and autumn 2013 (Appendix 4). Functional groups for the recorded taxa are listed in Appendix 3. In all wetlands changes in floristic composition through time were not consistent between elevations, as indicated by a significant interaction between elevation and time (Table 4).

Table 4: PERMANOVA results comparing the plant community through time and between elevations for each wetland.

Wetland	Factor	df	Pseudo-F	P
Angas and Bremer River Mouths	Time	9, 119	3.90	<0.001
	Elevation	1, 119	58.35	<0.001
	Time x Elevation	7, 119	3.16	<0.001
Dunn's Lagoon	Time	9, 599	10.15	<0.001
	Elevation	4, 599	79.53	<0.001
	Time x Elevation	36, 599	4.49	<0.001
Goolwa Channel Drive	Time	9, 269	7.82	<0.001
	Elevation	2, 269	73.46	<0.001
	Time x Elevation	18, 269	3.09	<0.001
Hunter's Creek	Time	9, 347	18.18	<0.001
	Elevation	1, 347	224.64	<0.001
	Time x Elevation	9, 347	8.13	<0.001
Loveday Bay	Time	9, 569	7.94	<0.001
	Elevation	4, 569	14.50	<0.001
	Time x Elevation	36, 569	2.55	<0.001
Milang	Time	9, 448	2.74	<0.001
	Elevation	3, 448	26.78	<0.001
	Time x Elevation	27, 448	1.41	0.002
Narrung	Time	9, 329	5.98	<0.001
	Elevation	3, 329	101.86	<0.001
	Time x Elevation	27, 329	2.99	<0.001
Poltalloch	Time	9, 239	14.39	<0.001
	Elevation	3, 239	26.27	<0.001
	Time x Elevation	27, 239	4.87	<0.001
Point Sturt	Time	9, 179	12.73	<0.001
	Elevation	2, 179	14.76	<0.001
	Time x Elevation	18, 179	3.00	<0.001
Teringie	Time	9, 479	5.59	<0.001
	Elevation	3, 479	39.90	<0.001
	Time x Elevation	27, 479	4.19	<0.001
Waltowa	Time	9, 239	12.03	<0.001
	Elevation	3, 239	57.02	<0.001
	Time x Elevation	21, 239	2.62	<0.001

3.1.1 Angas and Bremer River Mouths

A total of 36 taxa (including 25 exotics) were recorded at the mouths of the Angas and Bremer Rivers from spring 2008 to autumn 2013 (Appendix 4). On the edges of the streams, taxa from the terrestrial, amphibious and emergent functional groups were consistently present. In particular, the edges of both streams were dominated by *Pennisetum clandestinum* (terrestrial dry) throughout the survey period (Figure 6; Gehrig *et al.* 2011b). One exception was in autumn 2011 (nine months after inundation with lake water) when the submergent k-selected *Ceratophyllum demersum* was present. No significant indicator species were recorded for spring 2012 and autumn 2013 along the stream edges.

A greater functional diversity was recorded within channels throughout the survey period, where there was the general trend that in spring taxa from amphibious, emergent, floating and submergent k-selected functional groups were present as opposed to terrestrial, floodplain, amphibious and emergent functional groups recorded in autumn (Figure 6). However, there was the exception that following inundation with lake water in winter 2010, amphibious, emergent, submergent k-selected and floating taxa were recorded in autumn 2011, only amphibious and emergent taxa present in autumn 2012 and amphibious, emergent and floating taxa in autumn 2013 (Figure 6). For the channel, there was a significant increase in the floating taxa *Azolla filiculoides* in spring 2012 and *Lemna* spp. in autumn 2013.

3.1.2 Dunn's Lagoon

A total of 72 taxa (including 38 exotics) were recorded from Dunn's Lagoon from spring 2008 to autumn 2013 (Appendix 4). From spring 2008 until autumn 2010, the most abundant taxa were generally from floodplain and terrestrial functional groups, but by spring 2010 (when all but the highest elevation were inundated) there was a strong shift towards aquatic taxa (amphibious, emergent and submergent functional groups), particularly at the lower elevations (3 to 1, inclusive) (Figure 7). The highest elevation (5) had a significant increase of the terrestrial species *Distichlis distichophylla* in autumn 2013. At the next elevation (4), terrestrial taxa *Distichlis distichophylla* and *Plantago coronopus* were significantly more abundant in spring 2012. For elevation 3, *Azolla filiculoides*, *Juncus acutus*, *Myriophyllum caput-medusae* and *Typha domingensis* increased in abundance in spring 2012. At elevation 2, there was a significant increase in abundance of the native species *Myriophyllum caput-medusae* in spring 2012 and *Schoenoplectus validus*, *Azolla filiculoides*, and *Myriophyllum salsugineum* in autumn 2013. The lowest elevation (1) had a significant increase in the submergent *Myriophyllum caput-medusae* in spring 2012 and similarly in autumn 2013, *Myriophyllum salsugineum* and *Schoenoplectus validus* significantly increased.

3.1.3 Goolwa Channel Drive

A total of 32 taxa (including 18 exotics) were recorded at Goolwa Channel Drive Wetland from spring 2008 to autumn 2013 (Appendix 4). Throughout the survey period, the plant communities at the highest (3) and middle (2) elevations were dominated by terrestrial, emergent and amphibious taxa (Figure 8);

characterised by a native salt marsh community dominated by *Suaeda australis*, *Sarcocornia quinqueflora*, *Juncus kraussii*, *Frankenia pauciflora*, *Samolus repens*, *Distichlis distichophylla*, *Triglochin striatum* and *Schoenoplectus pungens* (Gehrig *et al.* 2011b). The highest elevation (3) had a significant increase in *Aster subulatus* in spring 2012, whereas *Phragmites australis* and *Paspalum distichum* significantly increased in abundance in autumn 2013. For the middle elevation (2) there were no significant changes in spring 2012, but a significant increase in the emergent *Phragmites australis* in autumn 2013. At the lowest elevation there were no significant indicators, and hence no significant changes, in spring 2012 and autumn 2013.

3.1.4 Hunter's Creek

A total of 38 taxa (including 16 exotics and one species listed as rare in South Australia) were recorded in Hunter's Creek from spring 2008 to autumn 2013 (Appendix 4). Across the survey period the plant community along the edges were predominantly amphibious, terrestrial and emergent taxa (Figure 9); although some submergent k-selected taxa were observed in the shallowly flooded margins (outside the surveyed quadrats) after the creek was inundated with lake water (winter 2010). In spring 2012, however, there was a significant increase in submergent (*Myriophyllum caput-medusae*, *Potamogeton crispus*, *Potamogeton pectinatus*, *Myriophyllum salsugineum*), floating (*Azolla filiculoides*), emergent (*Bolboschoenus caldwellii*) and amphibious (*Polypogon monspeliensis*, *Triglochin striatum*) taxa when edges were inundated. Similarly, there was an increase in abundance of submergent (*Ceratophyllum demersum*), floating (*Lemna* spp.) and terrestrial damp (*Paspalum distichum*) taxa in autumn 2013.

Prior to autumn 2010, submergent r-selected *Ruppia tuberosa* was present in the creek channel in spring 2008 and spring 2009, until amphibious halophytes *Suaeda australis* and *Sarcocornia quinqueflora* colonised the bed after drying out (Figure 9). After the creek was inundated in winter 2010, there was a significant increase in the abundance of submergent k-selected, amphibious, floating and terrestrial taxa (Figure 9) including *Potamogeton crispus*, *Potamogeton pectinatus* and *Myriophyllum caput-medusae* in spring 2012 and *Ceratophyllum demersum* and *Myriophyllum salsugineum* in autumn 2013.

3.1.5 Loveday Bay

A total of 53 taxa (including 32 exotics) were recorded from Loveday Bay from spring 2008 to autumn 2013 (Appendix 4). During the survey period, terrestrial, amphibious, floodplain and emergent taxa were present at the higher elevations (5 to 2, inclusive) (Figure 10), with the plant community largely dominated by *Sarcocornia quinqueflora* and *Suaeda australis*, which are both adapted to elevated salinities.

When the lowest elevation (1) was inundated (e.g. in spring or following inundation with lake water in winter 2010) submergent r-selected taxa: *Ruppia tuberosa* and *Lamprothamnium macropogon* were often present; albeit in significantly higher abundances in spring 2008 (Gehrig *et al.* 2011b). Otherwise at this elevation, emergent, amphibious and terrestrial taxa were present across the survey period (Figure 10). At the lowest elevation, there was a significant increase in filamentous green algae at the lowest elevation (1)

in spring 2012, but no changes in autumn 2013. At the next elevation (2) filamentous green algae and *Ruppia tuberosa* were also significant indicators in spring 2012, while *Chenopodium glaucum* was significant in autumn 2013. Similarly filamentous green and *Ruppia tuberosa* were significantly more abundant at elevation 3 in spring 2012, but *Chenopodium glaucum* and *Schoenoplectus pungens* were significant indicators in autumn 2013. There were no changes at the next elevation (4) in spring 2012, but *Typha domingensis* significantly increased in abundance in autumn 2013. At the highest elevation (5) there was a significant increase in *Juncus kraussii*, *Polypogon monspeliensis*, *Samolus repens* and *Trifolium* spp. in spring 2012, but in autumn 2013 *Aster subulatus* was the only significant indicator.

3.1.6 Milang

A total of 64 taxa (including 41 exotics; the largest number recorded of the wetlands surveyed) were recorded from Milang Wetland from spring 2008 to autumn 2013 (Appendix 4). Similar to Loveday Bay the plant community in Milang wetland was dominated by terrestrial, amphibious and some emergent taxa, across the entire survey (Figure 11), but changed seasonally with some winter annuals present during the spring surveys at all elevations. In addition, an area that was inundated seasonally with local runoff contained *Myriophyllum* sp. in spring 2008 and *Ruppia polycarpa* every spring during the survey period (abundance peaked in spring 2009) at some of the elevations (Figure 11; Gehrig *et al.* 2011b).

During the latest survey period (spring 2012 and autumn 2013) the plant community at the lowest elevation (4) and elevation 2 remained unchanged (i.e. no significant indicators), although at elevation 3 the floating species *Lemna* sp. significantly increased in abundance in spring 2012, while *Duma florulenta* and *Suaeda australis* increased in autumn 2013. At the highest elevation (1), there were no changes in spring 2012, but *Sarcocornia quinqueflora* significantly increased in abundance in autumn 2013.

3.1.7 Narrung

A total of 32 taxa (including 18 exotics) were recorded from Narrung Wetland from spring 2008 to autumn 2013 (Appendix 4). The plant communities at the highest elevations (4 to 2, inclusive) were a mix of terrestrial, amphibious and emergent taxa (Figure 12); in particular a native samphire community of *Sarcocornia quinqueflora*, *Suaeda australis* and *Frankenia pauciflora* (Gehrig *et al.* 2011b). The plant community remained unchanged at the highest (4) and lowest (1) elevations over spring 2012 and autumn 2013. At elevation 2, the plant community similarly remained unchanged in spring 2012, but *Bolboschoenus caldwellii* and *Paspalum distichum* were significant indicators in autumn 2013. Alternatively, *Polypogon monspeliensis* and *Sonchus oleraceus* significantly increased in abundance at elevation 3 in spring 2012, but there were no significant changes in autumn 2013.

3.1.8 Point Sturt

A total of 40 taxa (including 26 exotics) were recorded from Point Sturt Wetland from spring 2008 to autumn 2013 (Appendix 4). Similar to Dunns Lagoon, the wetland was generally dominated by terrestrial,

amphibious and floodplain taxa for the first two years of the survey period (Figure 13); however, following inundation of the wetland (winter 2010), the plant community shifted and exotic amphibious taxa dominated (Figure 13). Furthermore, in autumn 2011, emergent, submergent r-selected (*Ruppia polycarpa*) and some native floating taxa were also present (Figure 13; Gehrig *et al.* 2011b). During the latest survey period (spring 2012 to autumn 2013) there were no significant taxa, indicating the community remained unchanged, across all elevations.

3.1.9 Poltalloch

A total of 26 taxa (including 16 exotics) were recorded from Poltalloch Wetland from spring 2008 to autumn 2013 (Appendix 4). Prior to inundation with lake water (winter 2010), Poltalloch was dominated by terrestrial and amphibious taxa (Figure 14); largely salt tolerant taxa, with winter annuals present in the spring surveys. After inundation there was an increase in the abundance of amphibious taxa (especially at the higher elevations) that were later replaced by submergent r-selected taxa (e.g. *Ruppia tuberosa*) by autumn 2011. In the latest survey there were no significant indicators, except the submergent r-selected *Ruppia polycarpa* at elevation 2 in spring 2012.

3.1.10 Teringie

A total of 51 taxa (including 27 exotics) were recorded from Teringie Wetland from spring 2008 to autumn 2013 (Appendix 4). Prior to inundation (winter 2010), terrestrial, amphibious and floodplain salt tolerant taxa (*Sarcocornia quinqueflora*, *Suaeda australis*, *Distichlis distichophylla*) were present in the spring surveys (Figure 15; Gehrig *et al.* 2011b). Elevation 1 was inundated during spring 2010 from local runoff and the submergent r-selected species *Lamprothamnium macropogon*, *Ruppia tuberosa* and terrestrial *Spergularia brevifolia* had recruited (Figure 15; Gehrig *et al.* 2011b). Elevations 2 to 4 remained dominated by salt tolerant taxa for the corresponding period with winter annuals present in spring 2010. In spring 2012, *Azolla filiculoides*, *Lemna* spp. and *Plantago coronopus* significantly increased in abundance at the lowest elevation (1), but there were no changes in autumn 2013. The taxa *Azolla filiculoides*, *Lemna* spp. and *Samolus repens* also significantly increased in abundance at elevation 2 in spring 2012, but no significant changes were observed in autumn 2013. At elevation 3, *Lolium* spp. was a significant indicator in spring 2012 and *Paspalum distichum* significantly increased in autumn 2013. At the highest elevation (4) there were no significant changes in spring 2012, but there was a significant increase in *Paspalum distichum* in autumn 2013.

3.1.11 Waltowa

A total of 18 taxa (including 11 exotics) were recorded from Waltowa from spring 2008 to autumn 2013 (Appendix 4). The flow control structure on the wetland inlet was opened from May 2011 until April 2012. During that time the regulator was intermittently closed for short durations (days) to maintain water levels but between October 2011 and March 2012 there were difficulties in maintaining movement water into the wetland to compensate for the evaporative losses in the wetland due to choking of the inlet by

Phragmites australis (although this area was cleared in autumn 2012). As a result the wetland was often dry during this period because sufficient water could not be delivered consistently. The higher elevations (4 to 2, inclusive) were largely dominated by terrestrial, floodplain and amphibious taxa (Figure 16), which was primarily salt tolerant taxa such as, *Sarcocornia quinqueflora* and *Suaeda australis*. During the latest surveys (spring 2012 and autumn 2013) the plant community remained largely unchanged, except for a significant increase of *Reichardia tingitana* at the highest elevation (4) in spring 2012.

3.2 Lakeshores

A total of 106 taxa (63 exotics including one weed of national significance (*Tamarix aphylla*), three proclaimed pest plants in South Australia and one species listed as rare in South Australia) were recorded at shoreline sites in Lake Alexandrina, Lake Albert and Goolwa Channel from spring 2008 to autumn 2013. Lake Alexandrina had the highest species richness (87 taxa across 10 sites) followed by Goolwa Channel (70 taxa across 10 sites) with Lake Albert exhibiting the lowest species richness (58 taxa across 5 sites) (Appendix 5). Lake Albert had the highest proportion of exotics (58.62%) compared to Lake Alexandrina (49.54%) and Goolwa Channel (42.85%) (Appendix 5).

In each location (Lake Alexandrina, Lake Albert and Goolwa Channel) the plant community changed through time, was different between elevations and there was a significant interaction between the two factors (Table 5). This indicates that the plant community changes through time were not consistent between elevations for each location. Furthermore, the species (and hydrological processes) that caused the changes differed between locations.

Table 5: PERMANOVA results comparing the plant community through time and between elevations for Lake Alexandrina, Lake Albert and Goolwa Channel shorelines.

Elevation	Factor	df	F	P
Lake Alexandrina	Elevation	4, 459	4.45	<0.001
	Time	9, 459	18.05	<0.001
	Elevation × Time	36, 459	1.039	0.345
Lake Albert	Elevation	5, 779	24.60	<0.001
	Time	9, 779	39.06	<0.001
	Elevation × Time	43, 779	5.38	<0.001
Goolwa Channel	Elevation	4, 1259	16.35	<0.001
	Time	9, 1259	45.59	<0.001
	Elevation × Time	36, 1259	4.93	<0.001

3.2.1 Lake Alexandrina

Between spring 2008 and autumn 2010, the plant community in Lake Alexandrina was generally dominated by terrestrial taxa across all elevations (Figure 17). The following two years saw decreases in terrestrial taxa and a concomitant increase in emergent taxa, across all elevations (Figure 17). At the highest elevation (+0.8 m AHD) there was a significant increase in *Rumex bidens* in spring 2012 and *Lycopus australis* and *Typha domingensis* in autumn 2013. At the next elevation (+0.6 m AHD) significant

indicators for spring 2012 were *Azolla filiculoides* and *Hydrocotyle verticillata*, while *Persicaria lapathifolia* and *Typha domingensis* significantly increased in abundance in autumn 2013. At the +0.4 m AHD elevation, *Azolla filiculoides* significantly increased in abundance in spring 2012 while *Limosella australis* and *Typha domingensis* were significantly more abundant in autumn 2013. At +0.2 m AHD, there was a significant increase in *Azolla filiculoides* in spring 2012 and *Ludwigia peploides* and *Typha domingensis* were significant indicators in autumn 2013. At the 0 m AHD elevation, *Myriophyllum salsugineum* significantly increased in abundance in spring 2012, but there were no significant changes in autumn 2013. Lastly, there were no significant changes at the lowest elevation (-0.5 m AHD) in spring 2012 to autumn 2013.

3.2.2 Lake Albert

The plant community around the edge of Lake Albert, similar to Lake Alexandrina, was generally dominated by terrestrial taxa between spring 2008 and autumn 2010, with some floodplain and amphibious taxa also present (Figure 18). From spring 2010, amphibious and emergent taxa replaced the terrestrial taxa, particularly between elevations +0.4 and +0.6 m AHD (Figure 18). At elevation +0.2 AHD there was a marked increase in emergent taxa (Figure 18). Apart from a significant increase in the abundance of *Anagallis arvensis* at +0.8 m AHD in autumn 2013, there were no significant indicators for all remaining elevations, suggesting that the plant communities remained unchanged for the latest surveys in spring 2012 to autumn 2013.

3.2.3 Goolwa Channel

In Goolwa Channel, terrestrial, floodplain, amphibious and emergent taxa dominated the plant community at the higher elevations (0 to +0.8 m AHD, inclusive) between spring 2008 and autumn 2009, in contrast to the lowest elevation (-0.5 m AHD), which remained bare (Figure 19). When water levels rose in spring 2009, due to regulated flooding, some of the terrestrial taxa at these higher elevations were extirpated, although there was limited recruitment of terrestrial taxa above 0 m AHD when water levels fell in summer and autumn 2009-10 (Figure 19). There was little change in the abundances of the emergent fringing species (e.g. *Phragmites australis*) over the survey period in Goolwa Channel (Gehrig *et al.* 2011b). During the latest survey, at +0.8 m AHD, *Mentha* spp. significantly increased in abundance in spring 2012, but there were no significant changes in autumn 2013. At +0.6 m AHD the emergents *Berula erecta* and *Eleocharis acuta* were significant indicators in spring 2012, whereas *Typha domingensis* was significant in autumn 2013. At the next elevations (+0.4, +0.2 and 0 m AHD) there were no significant changes in spring 2012, but *Typha domingensis* significantly increased in abundance in autumn 2013. There were no significant indicator species for the lowest elevation (-0.5 m AHD) in spring 2012 and autumn 2013 suggesting there were no changes in the plant community across the latest survey period.

Table 6: Colour codes for vegetation functional groups.

Functional Group (colour codes)
Terrestrial
Floodplain
Amphibious
Floating
Emergent
Submergent r-selected
Submergent k-selected



**(A) Edge****(B) Channel**

Figure 6: Proportion of plant functional groups for edge (A) and channel (B) elevations of Angas and Bremer River Mouths from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).




Figure 7: Proportion of plant functional groups from highest (A: peg 5) to lowest elevation (E: peg 1) in Dunn’s Lagoon from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).






Figure 8: Proportion of plant functional groups from highest (A: peg 3) to lowest elevation (C: peg 1) in Goolwa Channel Drive from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).




Figure 9: Proportion of plant functional groups for edge (A) and channel (B) elevations of Hunter's Creek from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).







Figure 10: Proportion of plant functional groups across highest elevation (A: peg 4) to lowest elevation (E: peg 1) in Loveday Bay wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).




(A) peg 1



(B) peg 2




(C) peg 3




(D) peg 4


Figure 11: Proportion of plant functional groups across highest (A: peg 1) to lowest elevation (D: peg 4) in Milang Wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).




(A) Peg 4



(B) Peg 3




(C) Peg 2




(D) Peg 1


Figure 12: Proportion of plant functional groups across highest elevation (A: peg 4) to lowest elevation (D: peg 1) in Narrung wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).



(A) Peg 3



(B) Peg 2



(C) Peg 1

Figure 13: Proportion of plant functional groups across highest (A: peg 3) to lowest elevation (C: peg 1) in Point Sturt wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).







Figure 14: Proportion of plant functional groups across highest (A: peg 4) to lowest elevation (D: peg 1) in Poltalloch wetland from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).







Figure 15: Proportion of plant functional groups across highest (A: peg 4) to lowest elevation (D: peg 1) in Teringie from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).







Figure 16: Proportion of plant functional groups across highest elevation (A: peg 4) to lowest elevation (D: peg 1) in Waltowa from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).




Figure 17: Proportion of plant functional groups from highest elevation (+0.8 m AHD) to lowest elevation (-0.5 m AHD) in Lake Alexandrina from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).




Figure 18: Proportion of plant functional groups from highest elevation (+0.8 m AHD) to lowest elevation (-0.5 m AHD) in Lake Albert from spring 2008 to autumn 2013 (Functional group key is provided in Table 6).




Figure 19: Proportion of plant functional groups from highest elevation (+0.8 m AHD) to lowest elevation (-0.5 m AHD) in Goolwa Channel from spring 2008 to autumn 2013 (Functional group key is provided in Table 6

4 Discussion and management implications

During the most recent survey period (spring 2012 to autumn 2013) water levels and salinity in the Lake Alexandrina and Goolwa Channel remained similar to those recorded in the previous year (2011–2012) and salinity decreased in Lake Albert (although the decrease was probably not biologically significant for the plant species present). During the drought-induced draw down period (2007 to 2010), plant communities shifted towards an assemblage of terrestrial and floodplain taxa, but following restoration of water levels in the Lower Lakes in late August 2010 (and the subsequent reconnection of most wetlands) there has been an increase in the abundance and diversity of aquatic dependent taxa (e.g. submergent, amphibious, floating and emergent), suggesting the system may be beginning to recover and stabilise.

During 2012/13, water level management in the Lower Lakes involved two draw down and refilling cycles (between +0.4 and +0.8 m AHD) with the aim to reduce salinity in Lake Albert (Figure 2). Stable water levels have been identified as detrimental to aquatic plant communities, with a greater diversity of aquatic plants generally in systems with fluctuating water levels (e.g. Nielsen and Chick 1997). The managed water level fluctuations in the Lower Lakes periodically exposed the fringes of lakeshores and wetlands, which provides opportunities for species requiring exposure to germinate (e.g. *Persicaria lapathifolia*, *Ludwigia peploides*, *Juncus* spp.) (Nicol 2004). However, in the Lower Lakes there is probably limited opportunity for recruitment of species that require exposure to germinate due to most fringing areas being densely vegetated with emergent species such as *Typha domingensis* or *Phragmites australis*. Generally, shorelines that are not densely vegetated are subjected to wave action which can prevent seedlings from establishing (e.g. Foote and Kadlec 1988); nevertheless, water level fluctuations between +0.8 and +0.4 m AHD are recommended because areas of submergent vegetation are then maintained, the establishment of amphibious taxa in areas protected from wave action is facilitated (e.g. shorelines planted with *Schoenoplectus validus*) and salinity in Lake Albert is reduced.

4.1 Wetlands

Prior to 2007, the plant communities in the Lower Lakes wetlands were a diverse assemblage of submergent, amphibious, floating and emergent taxa (Renfrey *et al.* 1989; Holt *et al.* 2005; Nicol *et al.* 2006). However, from early 2007 to August 2010, wetlands were generally dominated by terrestrial taxa (Gehrig *et al.* 2010), which is typical of wetlands subjected to prolonged drawdown (e.g. Nicol 2010). Furthermore, 46 submergent, emergent and amphibious taxa that were recorded in the 2004 (Holt *et al.* 2005) and 2005 (Nicol *et al.* 2006) River Murray wetland baseline surveys for the Lower Lakes wetlands were not recorded between October 2008 and March 2010 (Gehrig *et al.* 2010). The only wetlands that supported wetland plant communities during this period were those that contained areas that received local runoff. These areas supported submergent (usually submergent r-selected), amphibious and emergent taxa communities (Marsland and Nicol 2009; Gehrig *et al.* 2010). For example, Goolwa

Channel Drive was dominated by emergent and amphibious taxa from 2008 to 2010, particularly at the lowest elevation (Gehrig *et al.* 2010). At the lowest elevation in Loveday Bay Wetland, submergent r-selected species (*Ruppia tuberosa* and *Lamprothamnium macropogon*) were present in spring 2008 (Gehrig *et al.* 2010). In Milang Wetland, submergent k-selected (*Myriophyllum salsugineum*), amphibious (*Triglochin striatum*, *Cyperus gymnocaulos*, *Juncus kraussii*, *Juncus usitatus*), submergent r-selected (*Ruppia polycarpa*) and emergent (*Bolboschoenus caldwellii*, *Eleocharis acuta*, *Phragmites australis*, *Triglochin procerum*) taxa were present (Gehrig *et al.* 2010). In the Angas and Bremer River mouths, submergent k-selected species (*Ceratophyllum demersum* and *Vallisneria australis*) were also present in spring surveys prior to August 2010. However, the water present in the channels of the Angas and Bremer mouths each spring was due to catchment inflows derived from rainfall in the eastern Mount Lofty Ranges and not local runoff. In addition, Narrung Wetland received environmental water via pumping, in spring 2009 and Paton and Bailey (2010) reported that the submergent species *Ruppia tuberosa*, *Ruppia polycarpa*, *Lepilaena cylindrocapa*, *Lepilaena preissii*, *Potamogeton pectinatus*, *Lamprothamnium* sp. and *Nitella* sp. recruited in response to watering.

In August 2010, water levels in the Lower Lakes returned to historical levels; however, hydrological restoration of wetlands occurred at different times (and in the case of Waltowa only just occurred following the latest survey in autumn 2012). Wetlands with good hydrological connections with the lakes (Dunns Lagoon, Angas River Mouth, Bremer River Mouth, Poltalloch, Hunters Creek, Point Sturt, Narrung and Goolwa Channel Drive) filled when water levels exceeded the sills on the inlets and were inundated from August 2010. Sandbars formed at the inlets of Loveday Bay (D. Chandler pers. comm.) and Teringie (D. Walker pers. comm.) wetlands, preventing the wetlands from filling until they were cleared in summer 2010-11. Significant areas of Milang Wetland are perched and primarily receive water from local runoff (A. Frears pers. comm.) although some low lying areas were inundated with lake water. Therefore, the response of the plant community to the return of historical lake levels differed between wetlands with respect to species and functional groups.

In the inundated areas of Dunn's Lagoon, the Angas and Bremer River mouths, Poltalloch, Hunters Creek, Goolwa Channel Drive and Point Sturt, the plant community showed a change from predominantly terrestrial taxa to amphibious, submergent, floating and emergent taxa. Whilst this change was not detected by indicator species analysis until autumn 2011, in some instances changes had occurred by the spring 2010 survey in the aforementioned wetlands but taxa from the aquatic functional groups were in lower abundances compared with the autumn 2011 survey. The abundance of aquatic taxa either increased or remained similar to levels recorded in the 2010/11 surveys, except in Poltalloch where abundances decreased. The reason for the decrease in submergent species in Poltalloch is unknown; however, this wetland was historically seasonal with inundation in spring and drying in autumn and summer. Most of the taxa recorded since then were submergent r-selected species (*Ruppia tuberosa*, *Ruppia polycarpa*); hence, the extended inundation may be unfavourable for this particular functional group.

Milang Wetland generally remained dominated by terrestrial taxa despite low lying areas being inundated with lake water (these areas were predominantly bare or sparsely vegetated by amphibious and emergent species). However in areas that were dominated by emergent and amphibious taxa throughout the study period, there appeared to be increased vigour and improved condition in spring 2010 and autumn 2011, which did not result in a significant increase in percentage cover (pers. obs.). It is unclear whether this observation was due to above average local rainfall during the last 24 months of the survey period (Bureau of Meteorology 2011a) or increased lake levels. In 2012/13, the plant community changed very little although there were slight increases in the abundance of salt tolerant taxa (e.g. *Sarcocornia quinqueflora* at the higher elevations).

Narrung Wetland is predominantly a saltmarsh; however, Holt *et al.* (2005) and Paton and Bailey (2010) reported that a diverse submergent plant community was present when inundated. Therefore, it is unknown why submergent plants were absent from the lower elevations during the spring 2010 and autumn 2011 surveys after the wetland was inundated with lake water. However, *Ruppia tuberosa* and charophytes had colonised the wetland bed at the lower elevations by spring 2011, and were still present in autumn 2012. In 2012/13, the vegetation communities changed very little, although there was an increase in emergent taxa and terrestrial weeds in the middle elevations.

Loveday Bay and Teringie wetlands were not inundated with lake water until summer 2010-11. The submergent r-selected species, *Ruppia tuberosa* and *Lamprothamnium macropogon* were present in spring 2010 at both sites due to low elevations being inundated by local runoff. Both wetlands dried over summer and submergent plants had not recruited by autumn 2011, despite inundation with lake water. Low elevations close to Lake Alexandrina in Teringie Wetland remained inundated throughout 2011/12 but were dominated by amphibious taxa and the diverse submergent plant community recorded in spring 2004 (Holt *et al.* 2005) had not re-established. Loveday Bay Wetland only fills when lake levels are very high and there are strong westerly winds and lake water only just reached the causeway in 2011/12, which was probably why submergent species had not recolonised the wetland. In 2012/13, filamentous algae was present in significant abundances at the lower elevations in spring. At the higher elevations there was a marked increase in the abundance of emergents and amphibious taxa.

Sites in Waltowa Wetland remained dry for the survey period, despite water levels in Lake Albert rising to a level that allowed water to enter the wetland and the inlet structure being open from May 2011 to April 2012. Despite a slight increase in the abundance of terrestrial weeds in spring 2012, the plant community in Waltowa Wetland remained largely unchanged in 2012/2013, continuing to be dominated by salt tolerant species and terrestrial species and will probably remain dominated by these functional groups in the future (unless extended inundation occurs).

4.2 Lakeshores

The changes in the plant community observed in Goolwa Channel, Lake Alexandrina and Lake Albert over the survey period were due to changes in water levels and salinity brought about by different management regimes for each location and the unregulated River Murray flow from August 2010. At the beginning of the survey period all three areas were connected, exhibiting similar water levels (Figure 2) and salinities (Figure 3) and in spring 2008 and autumn 2009, all elevations, at all locations were dominated by terrestrial taxa due to low water levels (*sensu* Nicol 2010). Construction of the Narrung Bund and Clayton Regulator resulted in fragmentation and the ability to manage water levels in Lake Albert and Goolwa Channel independently of Lake Alexandrina. The Narrung Bund was constructed to maintain water levels above -0.5 m AHD in Lake Albert by pumping water from Lake Alexandrina. The Clayton Regulator was constructed to maintain elevated water levels in Goolwa Channel, the lower Finnis River and lower Currency Creek by impounding flows from the Finnis River and Currency and Tookayerta Creeks and pumping from Lake Alexandrina. In addition, salinity in Goolwa Channel, whilst not a direct result of regulated flooding, (salinity was elevated in Goolwa Channel prior to regulator construction), was also higher than Lake Alexandrina (Figure 3). The unregulated flow (and subsequent breaching of the Clayton Regulator and Narrung Bund), resulted in reconnection, an increase in water levels (Figure 2) and a reduction in salinity (Figure 3) from August 2010.

From spring 2009 to autumn 2010, water levels ranged from -0.8 to -1.0 m AHD in Lake Alexandrina and 0 to -0.7 m AHD in Lake Albert (Figure 2). Hence, the plant community in both locations was dominated by terrestrial taxa, although fringing emergent species (predominately *Phragmites australis*) were present but disconnected from the lakes.

In contrast, water levels in Goolwa Channel over the same period, ranged from +0.75 m AHD in spring 2009, to -0.1 m AHD in autumn 2010 (Figure 2) due to the influence of the Clayton Regulator, inflows from the tributaries and pumping from Lake Alexandrina. The plant community during this period showed zonation in relation to water depth (*sensu* Spence 1982). At high elevations (+0.4 to +0.8 m AHD) the plant community was dominated by emergent and amphibious species such as *Phragmites australis*, *Duma florulenta*, *Typha domingensis* and *Calystegia sepium*. At intermediate elevations (0 to +0.4 m AHD) emergent species such as *Typha domingensis* and *Schoenoplectus validus*, that are adapted to deeper water, were common. At -0.5 m AHD only submergents (*Potamogeton pectinatus*, *Vallisneria australis*, *Ceratophyllum demersum* and *Myriophyllum salsugineum*) were present in quadrats. In addition, *Ruppia megacarpa* and *Ruppia polycarpa* (submergent species) were observed in low numbers outside of monitoring quadrats. Submergent taxa were not observed until autumn 2010, which was not unexpected as the spring 2009 survey was undertaken four weeks after pumping ceased and the majority of submergent taxa require longer than four weeks of inundation to germinate (Nicol and Ward 2010a; 2010b).

In autumn 2010, prior to the breaching of the Clayton Regulator, surface water electrical conductivity (EC) in some areas of Goolwa Channel exceeded 20,000 $\mu\text{S.cm}^{-1}$ (Figure 3); a level significantly higher than the reported tolerances of several of the emergent and submergent species present (Bailey *et al.* 2002). Extensive stands of *Typha domingensis* (maximum reported salinity tolerance of 8,000 $\mu\text{S.cm}^{-1}$), *Phragmites australis* (reported to show signs of severe stress at 15,000 $\mu\text{S.cm}^{-1}$) and *Schoenoplectus validus* (maximum reported salinity tolerance of 700 $\mu\text{S.cm}^{-1}$) (Bailey *et al.* 2002) were present in Goolwa Channel despite the elevated salinity. However, plants in Goolwa Channel had clearly regenerated from rhizomes, which support evidence from the seed bank assessment where *Typha domingensis* and *Schoenoplectus validus* did not germinate in salinities in excess of 5,000 $\mu\text{S.cm}^{-1}$ (Nicol and Ward 2010b). Nevertheless, seeds remained viable when subjected to salinities as high as 20,000 $\mu\text{S.cm}^{-1}$ for at least six weeks (Nicol and Ward 2010b). These results suggested that there are local salt tolerant ecotypes of the aforementioned species present in Goolwa Channel (and potentially throughout the Lower Lakes). However, little is known about the impacts of sub-lethal salinities, except that under elevated salinities these species are restricted to colonising new areas asexually.

In August 2010, River Murray inflows into Lake Alexandrina increased, water levels rose rapidly (Figure 2) and the Clayton Regulator and Narrung Bund were breached in September 2010, reconnecting the three locations. This resulted in similar water levels throughout the Lower Lakes (Figure 2); however, surface water EC in Lake Albert was significantly higher than Goolwa Channel and Lake Alexandrina (Figure 3). The terrestrial taxa that had recruited on the exposed sediment in Lakes Alexandrina and Albert were extirpated. Emergent and amphibious taxa increased in abundance from +0.8 to 0 m AHD in Lake Alexandrina but not in Lake Albert. In addition, species richness was lower in Lake Albert compared to Lake Alexandrina and Goolwa Channel. It is unlikely that the elevated surface water EC in Lake Albert is the cause of lower species richness, since levels have not exceeded thresholds for most species present throughout the system since reconnection (Bailey *et al.* 2002; Nicol and Ward 2010a; 2010b). The lower overall species richness is probably due to the dominance of a high energy shoreline in Lake Albert compared to Goolwa Channel and Lake Alexandrina where there is greater shoreline complexity and greater areas of protection from wave action.

The -0.5 m AHD elevation was generally devoid of plants in Lakes Albert and Alexandrina from spring 2010 onwards, except in one site near Clayton Bay in Lake Alexandria, where *Typha domingensis* and *Schoenoplectus validus* were present in low numbers. In Goolwa Channel, emergent taxa (especially *Schoenoplectus validus*) increased in abundance (probably due to lower surface water salinity) and there were significant changes to the submergent plant community between spring 2010 and autumn 2011. In spring 2010, the submergent plant community was dominated by *Potamogeton pectinatus*, which had colonised over 2,000 ha of Goolwa Channel, the lower Finniss River and lower Currency Creek at elevations between +0.4 m and -2.5 m AHD (Gehrig *et al.* 2011a). By autumn 2011, *Potamogeton pectinatus* had dramatically

decreased in distribution and abundance and areas previously dominated by this species were dominated by open water, *Myriophyllum salsgineum* and *Schoenoplectus validus* (Gehrig *et al.* 2011a).

The changes to the plant community in Goolwa Channel over the survey period can be attributed to changes in water quality (salinity and turbidity) and flow. *Potamogeton pectinatus* grows well in clear, slow flowing, saline water (at least 5‰ TDS, (approximately 7,500 $\mu\text{S}\cdot\text{cm}^{-1}$) (Sainty and Jacobs 2003); therefore, the conditions in Goolwa Channel between August 2009 and August 2010 were conducive for recruitment and spread of this species. Once established, *Potamogeton pectinatus* can colonise large areas rapidly by asexual reproduction (rhizomes and tubers) (Sainty and Jacobs 2003) and in Goolwa Channel large (almost monospecific) beds were present throughout the shallow water habitats (Gehrig *et al.* 2011a). In August 2010, fresh, turbid water replaced clear saline water and initially there was an increase in the abundance of *Potamogeton pectinatus* as there was sufficient photosynthetic tissue in the euphotic zone. However, as flows increased, plants were flattened and pushed out of the euphotic zone (and subsequently died) or were uprooted and washed into the Coorong. This provided an opportunity for species adapted to fresh turbid conditions (e.g. *Myriophyllum salsgineum*, *Ceratophyllum demersum*, *Potamogeton crispus*, *Vallisneria australis*) to colonise these areas.

Emergent taxa also responded to lower salinities in Goolwa Channel after September 2010 (Gehrig *et al.* 2011a). There has been an increase in abundance and extent of freshwater emergent species (*Typha domingensis*, *Phragmites australis* and *Schoenoplectus validus*) since the regulator was breached, which suggests that elevated salinity (whilst not lethal) reduced growth (Gehrig *et al.* 2011a).

Colonisation of submergent taxa in Goolwa Channel in response to regulated inundation and natural flooding provided evidence that the system is resilient and the aquatic plant community had the capacity to recover from low water levels. All submergent species observed in Goolwa Channel in autumn 2011 (except *Ceratophyllum demersum*) were present in the sediment seed bank (Nicol and Ward 2010b), which suggests that the seed bank is an important source of propagules for recolonisation of submergent taxa. Nevertheless, prior to reconnection with Lake Alexandrina, *Potamogeton crispus* was absent and *Ceratophyllum demersum* was restricted to the uppermost surveyed reaches of the Finnis River (adjacent to Wally's Landing). *Potamogeton crispus* and *Ceratophyllum demersum* are now widespread throughout Goolwa Channel (although not highly abundant) and are also present in Dunns Lagoon. Furthermore, *Myriophyllum caput-medusae* was present in the seed bank (Nicol and Ward 2010b) and historically present (J. Nicol pers. obs.) but absent from the extant vegetation since 2007. The absence of *Myriophyllum caput-medusae* was probably initially due to elevated salinity in Goolwa Channel, which exceeded the maximum reported salinity tolerance (Bailey *et al.* 2002) but presently is most likely due to lack of viable propagules and/or lack of dispersal.

Results from five years of monitoring show that allocation of sufficient water to maintain lake levels between +0.4 and +0.8 m AHD and provide periodic flushing to maintain low salinities produced the

most desirable outcomes with respect to aquatic plants. Construction of the regulators and pumping to maintain water levels should only be regarded as an emergency management action to mitigate acid sulfate soils. Nevertheless, regulated flooding resulted in recruitment of *Potamogeton pectinatus*, *Vallisneria australis* and *Myriophyllum salsugineum* and maintained emergent taxa, but salinities remained elevated throughout Goolwa Channel (water levels were not sufficiently high to inundate fringing vegetation in Lake Albert prior to reconnection). Elevated salinities resulted in reduced growth of emergent species and prevented or delayed germination (but not necessarily reduced seed viability) of emergent and submergent taxa (Nicol and Ward 2010b). This was supported by the distribution and abundance of three historically common freshwater submergent species (*Potamogeton crispus*, *Myriophyllum caput-medusae* and *Ceratophyllum demersum*) over the study period and the significant increase in the abundance of emergent species in autumn 2011. Furthermore, the dominance of *Potamogeton pectinatus* from March to November 2010 was probably due to regulated flooding and elevated salinity.

The salinity spikes in Goolwa Channel observed since April 2011 appeared to have little impact on the plant community. The salinity spikes were of short duration followed by flushing of freshwater and the maximum salinities were generally lower than the salinity experienced during autumn 2010.

4.3 The Living Murray Target V3

Whilst there has been a significant increase in the abundance of submergent, amphibious, floating and emergent species since water levels returned to historical levels, the abundances of species from these functional groups is now probably much lower than in 2004 (Holt *et al.* 2005) or 2005 (Nicol *et al.* 2006). For example, the diverse submergent plant communities that covered extensive areas in Clayton Bay, Dunn's Lagoon, Narrung, Milang (Holt *et al.* 2005), Loveday Bay, Hunters Creek, Point Sturt and Poltalloch (Nicol *et al.* 2006) had not re-established by autumn 2013. In addition, extensive areas of *Myriophyllum salsugineum* were present upstream of the Hindmarsh Island Bridge in shallow areas of Goolwa Channel prior to the drawdown of lake levels (pers. obs.) but were not present in autumn 2012. Furthermore, two species (*Lepilaena cylindrocarpa* and *Ranunculus trichophyllus*) recorded in the 2004 (Holt *et al.* 2005) and 2005 (Nicol *et al.* 2006) baseline surveys were not present in wetlands or lakeshores. However, extensive beds (468 ha) of *Myriophyllum salsugineum* had established in the lower Finnis River and lower Currency Creek, and the emergent plant community throughout Lake Alexandrina and Goolwa Channel showed signs of improved condition after water levels were reinstated. These plant communities were present throughout 2011/12 and whilst not measured directly probably increased in extent but still not to the level observed before the drought.

Therefore, TLM target V3 has not been met when compared to the plant community present prior to drawdown. However, all but two (*Ranunculus trichophyllus* and *Lepilaena cylindrocarpa*) of the 46 species recorded in 2004 and 2005, that were lost when water levels decreased, were recorded in the spring 2010 or autumn 2011 survey and were present in 2012/13. Results showed that the plant community shifted

from being dominated by terrestrial taxa to submergent, emergent, floating and amphibious taxa. Finally, most aquatic species are capable of rapid colonisation by asexual reproduction once mature (Grace 1993); therefore, it is likely that aquatic taxa will continue to increase in abundance providing water levels remain at current levels.

4.4 Further studies

Suggested further studies to improve the understanding of the vegetation dynamics of the Lower Lakes and impact of water levels and salinity include:

- Continue the condition monitoring program to gain an understanding of the medium- to long-term vegetation dynamics of the system and monitor recovery post hydrological restoration.
- Map large-scale plant communities in Goolwa Channel (*sensu* Gehrig *et al.* 2011a), expanding to key wetlands and lakeshore areas to complement the condition monitoring program and gain a better understanding of vegetation dynamics at the landscape scale.
- Undertake the *Melaleuca balmaturorum* component of the condition monitoring program in 2013/14.
- Define target or reference conditions for wetland/lakeshore/tributary communities.
- Investigate salinity tolerances of potential local ecotypes of key species.
- Investigate the effects of sub-lethal salinities on key species.
- Determine propagule longevity under different conditions (e.g. salinity, pH, soil moisture).
- Investigate current submergent plant propagule bank in key wetlands and Goolwa Channel.
- Investigate the relationships between plant communities and other biotic groups such as fish, birds and invertebrates.

5 References

- Abernethy, B. and Rutherford, I.D. (1998). Where along a river's length will vegetation most effectively stabilise stream banks? *Geomorphology* **23**: 55-75.
- Anderson, M.J. (2001). A new method for non-parametric analysis of variance. *Austral Ecology* **26**: 32-46.
- Anderson, M.J. and Ter Braak, C.J.F. (2003). Permutation tests for multi-factorial analysis of variance. *Journal of Statistical Computation and Simulation* **73**: 85-113.
- Bailey, P., Boon, P. and Morris, K. (2002). Salt sensitivity database. Land and Water Australia.
- Barker, W.R., Barker, R.M., Jessop, J.P. and Vonow, H.P. (2005). (Eds) Census of South Australian Vascular Plants (5.00 edn). (Botanic Gardens of Adelaide & State Herbarium: Adelaide)
- Bice, C., Wilson, P. and Ye, Q. (2008). Threatened fish populations in the Lower Lakes of the River Murray in spring 2007 and summer 2008. South Australian Research and Development Institute (Aquatic Sciences), F2008/000801-1, Adelaide.
- Blanch, S.J., Ganf, G.G. and Walker, K.F. (1999). Tolerance of riverine plants to flooding and exposure by water regime. *Regulated Rivers: Research and Management* **15**: 43-62.
- Blanch, S.J., Walker, K.F. and Ganf, G.G. (2000). Water regimes and littoral plants in four weir pools of the River Murray, Australia. *Regulated Rivers-Research & Management* **16**: 445-456.
- Brandle, R., Hammer, M., Wedderburn, S., Seaman, R., Noye, R. and Queale, L. (2002). A biological survey of the Murray Mouth Reserves, South Australia March 2002. Department for Environment and Heritage, Adelaide.
- Braun-Blanquet, J. (1932). 'Plant Sociology.' (McGraw-Hill: New York).
- Bray, J.R. and Curtis, J.T. (1957). An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs* **27**: 325-349.
- Brock, M.A. (1982). Biology of the salinity tolerant genus *Ruppia* L. in saline lakes in South Australia II. Population ecology and reproductive biology. *Aquatic Botany* **13**: 249-268.
- Brock, M.A. and Casanova, M.T. (1997). Plant life at the edge of wetlands: ecological responses to wetting and drying patterns. In 'Frontiers in Ecology: Building the Links'. (Eds Klomp, N. and Lunt, I.) pp. 181-192. (Elsevier Science: Oxford).

Brock, M.A., Smith, R.G.B. and Jarman, P.J. (1999). Drain it, dam it: alteration of water regime in shallow wetlands on the New England Tableland of New South Wales, Australia. *Wetlands Ecology and Management* **7**: 37-46.

Brownlow, M.D. (1997). Water Regime and the Aquatic Vegetation of Bool Lagoon, South Australia. PhD thesis, The University of Adelaide.

Bureau of Meteorology (2011a). Rainfall observations for Milang. <http://www.bom.gov.au/jsp/ncc/cdio/weatherData>

Camargo, A., Pezzato, M., Henry-Silva, G. and Assumpcao, A. (2006). Primary production of *Utricularia foliosa* L., *Egeria densa* Planchon and *Cabomba furcata* Schult & Schult.f from rivers of the coastal plain of the State of Sao Paulo, Brazil. *Hydrobiologia* **570**: 35-39.

Centre for Australian National Biodiversity Research and Council of Heads of Australasian Herbaria (2013). Australian Plant Census, IBIS database, <http://www.chah.gov.au/apc/index.html>.

Clarke, K.R. and Gorley, R.N. (2006). PRIMER version 6.1.12. (PRIMER-E Ltd: Plymouth).

Cronin, G., Lewis, W.M. and Schiehser, M.A. (2006). Influence of freshwater macrophytes on the littoral ecosystem structure and function of a young Colorado reservoir. *Aquatic Botany* **85**: 37-43.

Cunningham, G.M., Mulham, W.E., Milthorpe, P.L. and Leigh, J.H. (1981). 'Plants of Western New South Wales.' (New South Wales Government Printing Office: Sydney).

Dashorst, G.R.M. and Jessop, J.P. (1998). 'Plants of the Adelaide Plains and Hills.' (The Botanic Gardens of Adelaide and State Herbarium: Adelaide).

Declerck, S., Vandekerckhove, J., Johansson, L., Muylaert, K., Conde-Porcuna, J.M., Van der Gucht, K., Pérez-Martínez, C., Lauridsen, T., Schwenk, K., Zwart, G., Rommens, W., López-Ramos, J., Jeppesen, E., Vyverman, W., Brendonck, L. and De Meester, L. (2005). Multi-group biodiversity in shallow lakes along gradients of phosphorus and water plant cover. *Ecology* **86**: 1905–1915.

Department for Environment, Water and Natural Resources (2013a). Unpublished surface water electrical conductivity for Lake Alexandrina and Goolwa Channel from August 2008 to April 2013.

Department for Environment, Water and Natural Resources (2013b). Unpublished water levels for Lake Alexandrina and Goolwa Channel from August 2008 to April 2013.

dos Santos, A.M. and Esteves, F.D. (2002). Primary production and mortality of *Eleocharis interstincta* in response to water level fluctuations. *Aquatic Botany* **74**: 189-199.

- Dufrene, M. and Legendre, P. (1997). Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs* **67**: 345-366.
- Fitzpatrick, R., Grealish, G., Shand, P., Marvanek, S., Thomas, B., Creeper, N., Merry, R. and Raven, M. (2009a). Preliminary assessment of acid sulfate soil materials in Currency Creek, Finniss River, Tookayerta Creek and Black Swamp region, South Australia. CSIRO Land and Water, Adelaide.
- Fitzpatrick, R.W., Grealish, G., Shand, P., Simpson, S.L., Merry, R.H. and Raven, M.D. (2009b). Acid sulfate soil assessment in Finniss River, Currency Creek, Black Swamp and Goolwa Channel, South Australia. CSIRO Land and Water, 26/09, Adelaide.
- Foote, A.L. and Kadlec, J.A. (1988). Effects of wave energy on plant establishment in shallow lacustrine wetlands. *Journal of Freshwater Ecology* **4**: 523-532.
- Gehrig, S.L. and Nicol, J.M. (2010a). Aquatic and littoral vegetation monitoring of Goolwa Channel 2009-10. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2010/000383-1, Adelaide.
- Gehrig, S.L. and Nicol, J.M. (2010b). Aquatic and littoral vegetation of the Murray River downstream of Lock 1, the Lower Lakes, Murray Estuary and Coorong. A literature review. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2010/000297-1, Adelaide.
- Gehrig, S.L., Nicol, J.M. and Bucater, L. (2011a). Aquatic and littoral vegetation monitoring of Goolwa Channel 2009-11. South Australian Research and Development Institute (Aquatic Sciences), F2010/000383-2, Adelaide.
- Gehrig, S.L., Nicol, J.M. and Marsland, K.B. (2010). Lower Lakes aquatic and littoral vegetation condition monitoring 2009-10. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2010/000370-2, Adelaide.
- Gehrig, S.L., Nicol, J.M. and Marsland, K.B. (2011b). Lower Lakes vegetation condition monitoring – 2010/2011. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2009/000370-3, Adelaide.
- Gehrig, S.L., Nicol, J.M., Frahn, K.A. and Marsland, K.B. (2012). Lower Lakes vegetation condition monitoring – 2011/2012. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2010/000370-4, Adelaide.
- Grace, J.B. (1993). The adaptive significance of clonal reproduction in angiosperms: an aquatic perspective. *Aquatic Botany* **44**: 159-180.

- Grime, J.P. (1979). 'Plant Strategies and Vegetation Processes.' (John Wiley and Sons Ltd: Chichester).
- Hart, B.T., Lake, P.S., Webb, J.A. and Grace, M.R. (2003). Ecological risk to aquatic systems from salinity increases. *Australian Journal of Botany* **51**: 689-702.
- Heard, L. and Channon, B. (1997). Guide to a native vegetation survey using the biological survey of South Australia. Department of Environment and Natural Resources, Adelaide.
- Holt, M., Swingler, K., O'Donnell, E., Shirley, M., Lake, M., Conallin, A., Meredith, S., Ho, S., Prider, J., Poulsen, D., Richardson, S. and Cooling, M. (2005). River Murray Wetlands Baseline Survey. River Murray Catchment Water Management Board, Berri.
- James, K.R., Cant, B. and Ryan, T. (2003). Responses of freshwater biota to rising salinity levels and implications for saline water management: a review. *Australian Journal of Botany* **51**: 703-713.
- James, W.F., Best, E.P. and Barko, J.W. (2004). Sediment resuspension and light attenuation in Peoria Lake: can macrophytes improve water quality in this shallow system? *Hydrobiologia* **515**: 193-201.
- Jessop, J., Dashorst, G.R.M. and James, F.R. (2006). 'Grasses of South Australia. An illustrated guide to the native and naturalised species.' (Wakefield Press: Adelaide).
- Jessop, J.P. and Tolken, H.R. (1986). 'The Flora of South Australia.' (Government of South Australia Printer: Adelaide).
- Kingsford, R.T., Fairweather, P.G., Geddes, M.C., Lester, R.E., Sammut, J. and Walker, K.F. (2009). Engineering a crisis in a Ramsar Wetland: the Coorong, Lower Lakes and Murray Mouth, Australia. Australian Wetlands and Rivers Centre, The University of New South Wales, Sydney.
- Kingsford, R.T., Walker, K.F., Lester, R.E., Young, W.J., Fairweather, P.G., Sammut, J. and Geddes, M.C. (2011). A Ramsar wetland in crisis – the Coorong, Lower Lakes and Murray Mouth, Australia. *Marine and Freshwater Research* **62**: 255-265.
- Lang, P.J. and Kaeheneuhl, D.N. (2001). Plants of particular conservation significance in South Australia's agricultural regions. Department for Environment and Heriatage, Adelaide.
- Marsland, K. and Nicol, J. (2006). Current and proposed monitoring activities in relation to The Living Murray Lower Lakes, Coorong and Murray Mouth Icon Site Environmental Management Plan. SARDI Aquatic Sciences, SARDI Publication Number F2006/000081 P01, Adelaide.
- Marsland, K.B. and Nicol, J.M. (2009). Lower Lakes vegetation condition monitoring-2008/09. South Australian Research and Development Institute (Aquatic Sciences), F2009/000370-1, Adelaide.

- Maunsell Australia Pty Ltd. (2009). Lower Lakes, Coorong and Murray Mouth Icon Site condition monitoring plan. South Australian Murray-Darling Basin Natural Resources Management Board, Adelaide.
- McCune, B. and Mefford, M.J. (2006). PC-ORD. Multivariate Analysis of Ecological Data, Version 5.12. (MjM Software Design: Glenden Beach, Oregon, USA).
- Merry, R.H., Fitzpatrick, R.W., Barnett, E.J., Davies, P.J., Fotheringham, D.G., Thomas, B.P. and Hicks, W.S. (2003). South Australian inventory of acid sulfate soil risk (atlas). CSIRO Land and Water, Adelaide.
- Nicol, J. and Ward, R. (2010a). Seed bank assessment of Dunn's and Shadow's Lagoons. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication Number F2010/000594-1, Adelaide.
- Nicol, J. and Ward, R. (2010b). Seed bank assessment of Goolwa Channel, Lower Finniss River and Lower Currency Creek. South Australian Research and Development Institute (Aquatic Sciences), SARDI Publication No. F2010/000303-1, Adelaide.
- Nicol, J.M. (2004). Vegetation Dynamics of the Menindee Lakes with Reference to the Seed Bank. PhD thesis, The University of Adelaide.
- Nicol, J.M. (2010). Vegetation monitoring of River Murray Wetlands downstream of Lock 1. South Australian Research and Development Institute (Aquatic Sciences), F2009/000416-1, Adelaide.
- Nicol, J.M., Ganf, G.G. and Pelton, G.A. (2003). Seed banks of a southern Australian wetland: the influence of water regime on final species composition. *Plant Ecology* **168**: 191-205.
- Nicol, J.M. and Marsland, K.B. (2010). Vegetation monitoring in the Lower Finniss River, Currency Creek and Tookayerta Creek. South Australian Research and Development Institute (Aquatic Sciences), F2009/000674-1, Adelaide.
- Nicol, J.M., Weedon, J.T. and Doonan, A. (2006). Vegetation Surveys. In 'River Murray Wetlands Baseline Survey – 2005'. (Eds Simpson, D., Holt, M., Champion, T., Horan, A. and Shirley, M.). (South Australian Murray Darling Basin Natural Resources Management Board: Berri).
- Nielsen, D.L. and Brock, M.A. (2009). Modified water regime and salinity as a consequence of climate change: prospects for wetlands of Southern Australia. *Climate Change* **95**: 532-533.
- Nielsen, D.L., Brock, M.A., Rees, G.N. and Baldwin, D.S. (2003). Effects of increasing salinity on freshwater ecosystems in Australia. *Australian Journal of Botany* **51**: 655 – 665.

- Nielsen, D.L. and Chick, A.J. (1997). Flood-mediated changes in aquatic macrophyte community structure. *Marine and Freshwater Research* **48**: 153-157.
- Noges, T., Luup, H. and Feldmann, T. (2010). Primary production of aquatic macrophytes and their epiphytes in two shallow lakes (Peipsi and Vortsjarv) in Estonia. *Aquatic Ecology* **44**: 83-92.
- Papas, P. (2007). Effect of macrophytes on aquatic invertebrates – a literature review. Arthur Rylah Institute for Environmental Research, Department of Sustainability and Environment and Melbourne Water, Technical Report Series No. 158, Melbourne.
- Paton, D.C. and Bailey, C.P. (2010). Responses of aquatic vegetation to environmental watering of the Narrung wetland 2009-10. The University of Adelaide, Adelaide.
- Phillips, B. and Muller, K. (2006). Ecological character of the Coorong, Lakes Alexandrina and Albert Wetland of International Importance. Department for Environment and Heritage, Adelaide.
- Pinto, P., Morais, M., Ilheu, M. and Sandin, L. (2006). Relationships among biological elements (macrophytes, macroinvertebrates and ichthyofauna) for different core river types across Europe at two different spatial scales. *Hydrobiologia* **566**: 75-90.
- PIRSA Spatial Information Services (2009). Erosion mapping of the shorelines of Lakes Alexandina and Albert. Primary Industries and Resources South Australia (Spatial Information Services) and Goolwa to Wellington Local Action Planning Board Adelaide.
- Prescott, A. (1988). 'It's Blue with Five Petals. Wild Flowers of the Adelaide Region.' (Ann Prescott: Prospect, South Australia).
- Renfrey, A.P.C., Rea, N. and Ganf, G.G. (1989). The aquatic flora of Hindmarsh Island, South Australia. Department of Environment and Planning, Adelaide.
- Roberts, J. and Marston, F. (2000). Water Regime of Wetland and Floodplain Plants in the Murray-Darling Basin. CSIRO Land and Water, 30-00, Canberra.
- Romanowski, N. (1998). 'Aquatic and Wetland Plants. A Field Guide for Non-tropical Australia.' (University of New South Wales Press: Sydney).
- Sainty, G.R. and Jacobs, S.W.L. (1981). 'Water Plants of New South Wales.' (Water Resources Commission New South Wales: Sydney).
- Sainty, G.R. and Jacobs, S.W.L. (2003). 'Waterplants in Australia.' (Sainty and Associates: Darlinghurst, N.S.W., Australia).

- Spence, D.H.N. (1982). The zonation of plants in freshwater lakes. *Advances in Ecological Research* **12**: 37-125.
- Van Dijk, I.J.M., Beck, H.E., Crosbie, R.S., de Jeu, R.A.M., Liu, Y.Y., Podger, G.M., Timbal, B. and Viney, N.R. (2013). The Millenium Drought in southeast Australia (2001–2009): natural and human causes and implications for water resources, ecosystems, economy, and society. *Water Resources Research* **49**: 1040-1057.
- Webster, I.T., Parslow, J.S., Grayson, R.B., Molloy, R.P., Andrewartha, J., Sakov, P., Tan, K.S., Walker, S.J. and Wallace, B.B. (2001). Gippsland Lakes environmental study: assessing options for improving water quality and ecological function. Gippsland Coastal Board, Melbourne.
- Wedderburn, S.D., Walker, K.F. and Zampatti, B.P. (2007). Habitat separation of *Craterocephalus* (Atherinidae) species and populations in off-channel areas of the lower River Murray, Australia. *Ecology of Freshwater Fish* **16**: 442–449.

6 Appendices

Appendix 1: GPS coordinates (UTM format, map datum WGS84) for lakeshore and wetland understory vegetation monitoring sites.

Site	Easting	Northing	Site type
Bremer Mouth Lakeshore	323061	6081991	lakeshore
Brown Beach 1	350172	6052777	lakeshore
Brown Beach 2	350287	6053158	lakeshore
Clayton Bay	311301	6070626	lakeshore
Currency Creek 3	296772	6074222	lakeshore
Currency Creek 4	301013	6071800	lakeshore
Goolwa North	303330	6070156	lakeshore
Goolwa South	300490	6066366	lakeshore
Hindmarsh Island Bridge 01	299670	6068521	lakeshore
Hindmarsh Island Bridge 02	299695	6068616	lakeshore
Lake Reserve Rd	339298	6089987	lakeshore
Loveday Bay	329431	6058407	lakeshore
Loveday Bay Lakeshore	326621	6061647	lakeshore
Lower Finnis 02	305131	6076401	lakeshore
Milang	315964	6079870	lakeshore
Milang Lakeshore	316081	6079746	lakeshore
Pt Sturt Lakeshore	322811	6069643	lakeshore
Pt Sturt Water Reserve	317673	6070784	lakeshore
Teringie Lakeshore	327461	6066887	lakeshore
Upstream of Clayton Regulator	312281	6069151	lakeshore
Wally's Landing	303066	6079631	lakeshore
Warrenge 1	347722	6049163	lakeshore
Lower Finnis 03	305131	6072406	lakeshore
Narrung Lakeshore	333762	6069807	lakeshore
Nurra Nurra	341786	6063837	lakeshore
Warrenge 2	348487	6049133	lakeshore
Angas Mouth	318391	6081206	wetland
Bremer Mouth	323056	6082019	wetland
Dunns Lagoon	312417	6070300	wetland
Goolwa Channel Drive	307024	6064437	wetland
Hunters Creek	308219	6065526	wetland
Poltalloch	343248	6071554	wetland
Pt Sturt	322778	6069794	wetland
Teringie	327334	6065286	wetland
Waltowa	353908	6057756	wetland
Narrung	334542	6068744	wetland

Appendix 2: Current nomenclature changes Centre for Australian National Biodiversity Research and Council of Heads of Australasian Herbaria (2013).

Common Name	Previous scientific name		Revised scientific name	
	Genus	Species	Genus	Species
Blackseed samphire	<i>Halosarcia</i>	<i>pergranulata</i>	<i>Tecticornia</i>	<i>pergranulata</i>
Lignum	<i>Muehlenbeckia</i>	<i>florulenta</i>	<i>Duma</i>	<i>florulenta</i>
Weedy cudweed	<i>Pseudognaphalium</i>	<i>luteoalbum</i>	<i>Helichrysum</i>	<i>luteoalbum</i>
Brome grass	<i>Bromus</i>	<i>uniloides</i>	<i>Bromus</i>	<i>catharticus</i>
Soft Brome	<i>Bromus</i>	<i>mollis</i>	<i>Bromus</i>	<i>hordeaceus</i> ssp. <i>hordeaceus</i>
Angled Lobelia	<i>Lobelia</i>	<i>alata</i>	<i>Lobelia</i>	<i>anceps</i>
Native Picris	<i>Picris</i>	<i>hieracoides</i>	<i>Picris</i>	<i>angustifolia</i> ssp. <i>angustifolia</i>
Bog yellowcress	<i>Rorippa</i>	<i>islandica</i>	<i>Rorippa</i>	<i>palustris</i>
Ribbonweed	<i>Vallisneria</i>	<i>spiralis</i> var. <i>americana</i>	<i>Vallisneria</i>	<i>australis</i>
Thread-leaved water-crowfoot	<i>Batrachium</i>	<i>trichophyllum</i>	<i>Ranunculus</i>	<i>trichophyllum</i>
Prickly saltwort, tumbleweed	<i>Salsola</i>	<i>kali</i>	<i>Salsola</i>	<i>australis</i>
Short stem sand-spurrey	<i>Spergularia</i>	<i>marina</i>	<i>Spergularia</i>	<i>brevifolia</i>*

Appendix 3: Species list, functional classification (Gehrig and Nicol 2010b), life history strategy and conservation status (state conservation status from listings in Barker *et al.* (2005) and regional conservation status from listings in Lang and Kraehenuehl (2001) from all sites and survey dates (*denotes exotic taxon, **denotes proclaimed pest plant in South Australia, ***denotes weed of national significance # denotes listed as rare in South Australia).

Taxon	Functional Group	Life history strategy	Status and Comments
<i>Acacia myrtifolia</i>	Terrestrial dry	Perennial	Native
<i>Agapanthus praecox</i> *	Terrestrial dry	Perennial	Exotic
<i>Anagallis arvensis</i> *	Terrestrial damp	Annual	Exotic
<i>Apium graveolens</i> *	Terrestrial damp	Annual	Exotic
<i>Arctotheca calendula</i> *	Terrestrial dry	Annual	Exotic
<i>Asparagus asparagoides</i> ***	Terrestrial dry	Perennial	Exotic
<i>Asparagus officinalis</i> *	Terrestrial dry	Perennial	Exotic
<i>Asphodelus fistulosus</i> **	Terrestrial dry	Perennial	Exotic
<i>Aster subulatus</i> *	Terrestrial damp	Annual	Exotic
<i>Atriplex prostrata</i> *	Terrestrial damp	Perennial	Exotic
<i>Atriplex semibaccata</i>	Terrestrial dry	Perennial	Native-Listed as Uncommon in the Murray Region
<i>Atriplex stipitata</i>	Terrestrial dry	Perennial	Native
<i>Atriplex suberecta</i>	Floodplain	Perennial	Native
<i>Avena</i> spp.*	Terrestrial dry	Annual	Exotic- <i>Avena</i> spp. is comprised of <i>Avena barbata</i> and <i>Avena fatua</i>
<i>Azolla filiculoides</i>	Floating	Perennial	Native
<i>Berula erecta</i>	Emergent	Perennial	Native
<i>Bolboschoenus caldwellii</i>	Emergent	Perennial	Native
<i>Brassica rapa</i> *	Terrestrial dry	Annual	Exotic
<i>Brassica tournifortii</i> *	Terrestrial dry	Annual	Exotic
<i>Briza minor</i> *	Terrestrial dry	Annual	Exotic
<i>Bromus catharticus</i> *	Terrestrial dry	Annual	Exotic
<i>Bromus diandrus</i> *	Terrestrial dry	Annual	Exotic
<i>Bromus hordeaceus</i> ssp. <i>hordeaceus</i> *	Terrestrial dry	Annual	Exotic
<i>Calystegia sepium</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native-Listed as Uncommon in the Murray and Southern Lofty Regions
<i>Carex fascicularis</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Carpobrotus rossii</i>	Terrestrial dry	Perennial	Native
<i>Centaurea calcitrapa</i> *	Terrestrial damp	Annual	Exotic
<i>Centaureum tenuiflorum</i> *	Terrestrial damp	Annual	Exotic
<i>Ceratophyllum demersum</i> #	Submergent (k-selected)	Perennial	Native-Listed as Rare in South Australia
<i>Chara</i> spp.	Submergent (r-selected)	Annual	Native
<i>Chenopodium album</i> *	Terrestrial damp	Annual	Exotic
<i>Chenopodium glaucum</i> *	Terrestrial damp	Annual	Exotic
<i>Chenopodium nitrariaceum</i>	Terrestrial dry	Perennial	Native
<i>Conyza bonariensis</i> *	Terrestrial damp	Annual	Exotic
<i>Cotula coronopifolia</i> *	Amphibious fluctuation responder-plastic	Perennial	Exotic
<i>Crinum</i> sp.*	Terrestrial dry	Perennial	Exotic-garden escapee not in any of the identification keys and could not be identified to species
<i>Cyperus exaltatus</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Cyperus gymnocaulos</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Dianella revoluta</i>	Terrestrial dry	Perennial	Native
<i>Disphyma crassifolium</i>	Terrestrial dry	Perennial	Native
<i>Distichlis distichophylla</i>	Terrestrial damp	Perennial	Native-Listed as Uncommon in the Murray Region
<i>Duma florulenta</i>	Amphibious fluctuation	Perennial	Native

Taxon	Functional Group	Life history strategy	Status and Comments
	tolerator-woody		
<i>Echinochloa crus-galli</i> *	Terrestrial damp	Annual	Exotic
<i>Ehrharta longiflora</i> *	Terrestrial damp	Annual	Exotic
<i>Einadia nutans</i>	Terrestrial dry	Perennial	Native
<i>Eleocharis acuta</i>	Emergent	Perennial	Native
<i>Enchylaena tomentosa</i>	Terrestrial dry	Perennial	Native
<i>Epilobium pallidiflorum</i>	Terrestrial damp	Perennial	Native-Listed as Uncertain in the Murray Region and uncommon in the Southern Lofty Region
<i>Eragrostis australasica</i>	Floodplain	Perennial	Native
<i>Eragrostis curvula</i> **	Terrestrial damp	Annual	Exotic-Proclaimed pest plant in SA
<i>Eragrostis</i> sp.	Terrestrial damp	Annual	Native-could not identify to species
<i>Erodium cicutarium</i> *	Terrestrial dry	Annual	Exotic
<i>Euphorbia terracina</i> *	Terrestrial dry	Annual	Exotic-Proclaimed pest plant in SA
<i>Ficinia nodosa</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Foeniculum vulgare</i> *	Terrestrial damp	Annual	Exotic
<i>Frankenia pauciflora</i>	Terrestrial dry	Perennial	Native
<i>Fumaria bastardii</i> *	Terrestrial damp	Annual	Exotic
<i>Gahnia filum</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native-Listed as Rare in the Murray and Southern Lofty Regions
<i>Glyceria australis</i>	Emergent	Perennial	Native
<i>Helichrysum luteo-album</i>	Floodplain	Annual	Native
<i>Heliotropium europaeum</i> *	Floodplain	Annual	Exotic
<i>Holcus lanatus</i> *	Terrestrial damp	Annual	Exotic
<i>Hordeum vulgare</i> *	Terrestrial dry	Annual	Exotic
<i>Hydrocotyle verticillata</i>	Amphibious fluctuation responder-plastic	Perennial	Native-Listed as Uncertain in the Southern Lofty Region
<i>Hypochoeris glabra</i> *	Terrestrial dry	Annual	Exotic
<i>Hypochoeris radicata</i> *	Terrestrial dry	Annual	Exotic
<i>Iris</i> spp.*	Terrestrial dry	Perennial	Exotic
<i>Isolepis platycarpa</i>	Amphibious fluctuation tolerator-low growing	Perennial	Native
<i>Isolepis producta</i>	Amphibious fluctuation tolerator-low growing	Perennial	Native
<i>Juncus acutus</i> *	Amphibious fluctuation tolerator-emergent	Perennial	Exotic
<i>Juncus kraussii</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Juncus subsecundus</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Juncus usitatus</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Lachnagrostis filiformis</i>	Floodplain	Annual	Native
<i>Lactuca saligna</i> *	Terrestrial dry	Annual	Exotic
<i>Lactuca serriola</i> *	Terrestrial dry	Annual	Exotic
<i>Lagurus ovatus</i> *	Terrestrial dry	Annual	Exotic
<i>Lamprothamnium macropogon</i>	Submergent r-selected	Annual	Native
<i>Lemna</i> spp.	Floating	Perennial	Native
<i>Limosella australis</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Lobelia anceps</i>	Terrestrial damp	Perennial	Native
<i>Lolium</i> spp.*	Terrestrial dry	Annual	Exotic- <i>Lolium</i> spp. comprises of <i>Lolium perenne</i> and <i>Lolium rigidum</i>
<i>Ludwigia peploides</i> ssp. <i>montevidensis</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Lupinus cosentinii</i> *	Terrestrial dry	Annual	Exotic

Taxon	Functional Group	Life history strategy	Status and Comments
<i>Lycium ferocissimum</i> ***	Terrestrial dry	Perennial	Exotic-Proclaimed pest plant in SA
<i>Lycopus australis</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native-Listed as Rare in the Murray Region
<i>Lythrum hyssopifolia</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Lythrum salicaria</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Malva parviflora</i> *	Terrestrial dry	Annual	Exotic
<i>Marrubium vulgare</i> **	Terrestrial dry	Annual	Exotic
<i>Medicago</i> spp.*	Terrestrial dry	Annual	Exotic- <i>Medicago</i> spp. comprises of <i>Medicago polymorpha</i> , <i>Medicago truncatula</i> and <i>Medicago minima</i>
<i>Melaleuca halmaturorum</i>	Amphibious fluctuation tolerator-woody	Perennial	Native
<i>Melilotus indica</i> *	Terrestrial dry	Annual	Exotic
<i>Mentha australis</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Mentha</i> spp.*	Amphibious fluctuation tolerator-emergent	Perennial	Exotic- <i>Mentha</i> spp. comprises of <i>Mentha piperita</i> , <i>Mentha pulegium</i> and <i>Mentha spicata</i>
<i>Mimulus repens</i>	Amphibious fluctuation tolerator-low growing	Perennial	Native
<i>Muehlenbeckia gunnii</i>	Terrestrial dry	Perennial	Native
<i>Myriophyllum caput-medusae</i>	Submergent k-selected	Perennial	Native
<i>Myriophyllum salsugineum</i>	Submergent k-selected	Perennial	Native-Listed as Uncertain in the Southern Lofty Region
<i>Myriophyllum</i> sp.	Submergent k-selected	Perennial	Native
<i>Onopordum acanthium</i> *	Terrestrial damp	Annual	Exotic
<i>Oxalis pes-caprae</i> **	Terrestrial dry	Annual	Exotic-Proclaimed pest plant in SA
<i>Paspalum distichum</i> *	Terrestrial damp	Perennial	Exotic
<i>Pennisetum clandestinum</i> *	Terrestrial dry	Perennial	Exotic
<i>Persicaria lapathifolia</i>	Amphibious fluctuation responder-plastic	Perennial	Native
<i>Phalaris arundinacea</i> *	Amphibious fluctuation tolerator-emergent	Perennial	Exotic
<i>Phragmites australis</i>	Emergent	Perennial	Native
<i>Phyla canescens</i> *	Amphibious fluctuation tolerator-low growing	Perennial	Exotic
<i>Picris angustifolia</i> ssp. <i>angustifolia</i>	Terrestrial dry	Annual	Native
<i>Plantago coronopus</i> *	Terrestrial dry	Annual	Exotic
<i>Plantago lanceolata</i> *	Terrestrial dry	Annual	Exotic
<i>Plantago major</i> *	Terrestrial dry	Annual	Exotic
<i>Poa annua</i> *	Terrestrial damp	Annual	Exotic
<i>Polygonum aviculare</i> *	Terrestrial dry	Perennial	Exotic
<i>Polypogon monspeliensis</i> *	Amphibious fluctuation tolerator-emergent	Annual	Exotic
<i>Potamogeton crispus</i>	Submergent k-selected	Perennial	Native
<i>Potamogeton pectinatus</i>	Submergent k-selected	Perennial	Native
<i>Puccinellia</i> sp.*	Terrestrial damp	Annual	Exotic-could not be identified to species but was not <i>Puccinellia stricta</i> or <i>Puccinellia perluxa</i>
<i>Ranunculus trichophyllus</i> *	Submergent (r-selected)	Annual	Exotic
<i>Ranunculus trilobus</i> *	Amphibious fluctuation tolerator-emergent	Annual	Exotic
<i>Reichardia tingitana</i> *	Terrestrial dry	Annual	Exotic
<i>Rhagodia spinescens</i>	Terrestrial dry	Perennial	Native
<i>Rorippa nasturtium-aquaticum</i> *	Amphibious fluctuation responder-plastic	Annual	Exotic
<i>Rorippa palustris</i> *	Floodplain	Annual	Exotic
<i>Ruppia megacarpa</i>	Submergent k-selected	Perennial	Native
<i>Ruppia polycarpa</i>	Submergent r-selected	Annual	Native

Taxon	Functional Group	Life history strategy	Status and Comments
<i>Ruppia tuberosa</i>	Submergent r-selected	Annual	Native
<i>Salix babylonica</i> *	Emergent	Perennial	Exotic
<i>Salsola australis</i>	Terrestrial dry	Perennial	Native
<i>Samolus repens</i>	Terrestrial damp	Perennial	Native- Listed as Rare in the Murray Region and Uncommon the Southern Lofty Region
<i>Sarcocornia quinqueflora</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Scabiosa atropurpurea</i> *	Terrestrial dry	Annual	Exotic
<i>Schoenoplectus pungens</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native-Listed as Rare in the Southern Lofty Region
<i>Schoenoplectus validus</i>	Emergent	Perennial	Native
<i>Sclerolaena blackiana</i>	Terrestrial dry	Perennial	Native-Listed as Rare in SA
<i>Senecio cunninghamii</i>	Floodplain	Perennial	Native
<i>Senecio pterophorus</i> *	Terrestrial dry	Annual	Exotic
<i>Senecio runcinifolius</i>	Floodplain	Perennial	Native-Listed as Uncommon in the Murray Region
<i>Silybum marianum</i> **	Terrestrial damp	Annual	Exotic-Proclaimed pest plant in SA
<i>Solanum nigrum</i> *	Terrestrial damp	Annual	Exotic
<i>Sonchus asper</i> *	Terrestrial damp	Annual	Exotic
<i>Sonchus oleraceus</i> *	Terrestrial damp	Annual	Exotic
<i>Spergularia brevifolia</i> *	Terrestrial damp	Annual	Exotic
<i>Suaeda australis</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Tamarix aphylla</i> ***	Terrestrial dry	Perennial	Exotic
<i>Tecticornia pergranulata</i>	Amphibious fluctuation tolerator-emergent	Perennial	Native
<i>Trifolium</i> spp.*	Terrestrial dry	Annual	Exotic- <i>Trifolium</i> spp. comprises of <i>Trifolium angustifolium</i> , <i>Trifolium arvense</i> , <i>Trifolium repens</i> and <i>Trifolium subterraneum</i>
<i>Triglochin procerum</i>	Emergent	Perennial	Native-Listed as Uncommon in the Southern Lofty Region
<i>Triglochin striatum</i>	Amphibious fluctuation tolerator-low growing	Perennial	Native
<i>Triticum</i> sp.*	Terrestrial dry	Annual	Exotic-could not be identified to species
<i>Typha domingensis</i>	Emergent	Perennial	Native
<i>Urtica urens</i> *	Terrestrial damp	Annual	Exotic
<i>Vallisneria australis</i>	Submergent k-selected	Perennial	Native-Listed as Uncommon in the Murray Region and Threatened in the Southern Lofty Region
<i>Vicia sativa</i> *	Terrestrial dry	Annual	Exotic
<i>Wilsonia rotundifolia</i>	Terrestrial damp	Perennial	Native

Appendix 4: Taxa present (P) in each wetland from spring 2008 to autumn 2013 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia).

Species	Angas and Bremer Mouths	Dunns Lagoon	Goolwa Channel Drive	Hunters Creek	Loveday Bay	Milang	Narrung	Point Sturt	Poltalloch	Teringie	Waltowa
<i>Agapanthus</i> sp.*						P					
<i>Asparagus asparagoides</i> ***						P					
<i>Asphodelus fistulosus</i> **						P					
<i>Aster subulatus</i> *	P	P	P	P	P	P	P	P		P	
<i>Atriplex prostrata</i> *	P	P	P	P	P	P	P	P			
<i>Atriplex semibaccata</i>							P			P	
<i>Atriplex</i> spp.								P		P	
<i>Atriplex stipitata</i>					P						
<i>Atriplex suberecta</i>								P		P	
<i>Avena</i> spp.*	P	P				P		P		P	
<i>Azolla filiculoides</i>	P	P		P						P	
<i>Berula erecta</i>	P										
<i>Bolboschoenus caldwellii</i>			P	P	P	P	P	P	P		
<i>Brassica rapa</i> *		P									
<i>Brassica tournifortii</i> *		P				P				P	
<i>Bromus catharticus</i> *											P
<i>Bromus diandrus</i> *		P			P	P		P		P	
<i>Bromus hordeaceus</i> ssp. <i>hordeaceus</i> *		P			P	P		P		P	
<i>Calystegia sepium</i>		P									
<i>Carpobrotus rossii</i>							P			P	P
<i>Centaurea calcitrapa</i> *	P		P		P				P		
<i>Ceratophyllum demersum</i> #	P	P		P							
<i>Chara</i> spp.			P				P	P	P		
<i>Chenopodium album</i> *	P	P				P		P		P	
<i>Chenopodium glaucum</i> *		P			P			P	P		

Species	Angas and Bremer Mouths	Dunns Lagoon	Goolwa Channel Drive	Hunters Creek	Loveday Bay	Milang	Narrung	Point Sturt	Poltalloch	Teringie	Waltowa
<i>Conyza bonariensis</i> *	P	P	P		P	P					
<i>Cotula coronopifolia</i> *	P	P	P	P	P	P	P	P	P	P	
<i>Crinum</i> sp.						P					
<i>Cyperus exaltatus</i>	P										
<i>Cyperus gymnocaulos</i>		P			P	P			P	P	
<i>Dianella revoluta</i>		P									
<i>Disphyma crassifolium</i> ssp. <i>clavellatum</i>							P			P	P
<i>Distichlis distichophylla</i>		P	P	P	P	P	P	P	P	P	
<i>Echinochloa crus-galli</i> *				P							
<i>Ehrharta longiflora</i> *						P					
<i>Einadia nutans</i>								P			
<i>Eleocharis acuta</i>				P	P	P					
<i>Enchylaena tomentosa</i>		P				P		P		P	
<i>Eragrostis curvula</i> **		P		P	P	P	P	P	P	P	P
<i>Eragrostis</i> sp.		P				P	P	P		P	
<i>Erodium cicutarium</i> *										P	
<i>Ficinia nodosa</i>		P	P	P					P		
<i>Frankenia pauciflora</i>			P		P	P	P			P	
<i>Fumaria bastardii</i> *						P					
<i>Gahnia filum</i>		P									
<i>Glyceria australis</i>			P							P	P
<i>Helichrysum luteoalbum</i>		P			P						
<i>Heliotropium europaeum</i> *										P	
<i>Holcus lanatus</i> *						P					
<i>Hordeum vulgare</i> *		P			P	P	P	P	P	P	P
<i>Hydrocotyle verticillata</i>		P									
<i>Hypochaeris glabra</i> *		P			P						

Species	Angas and Bremer Mouths	Dunns Lagoon	Goolwa Channel Drive	Hunters Creek	Loveday Bay	Milang	Narrung	Point Sturt	Poltalloch	Teringie	Waltowa
<i>Hypochaeris radicata</i> *	P	P									
<i>Iris</i> sp.*	P										
<i>Isolepis platycarpa</i>		P				P					
<i>Isolepis producta</i>				P	P	P					
<i>Juncus acutus</i> *		P									
<i>Juncus kraussii</i>		P	P	P	P	P					
<i>Juncus subsecundus</i>		P		P	P	P					
<i>Juncus usitatus</i>						P					
<i>Lachnagrostis filiformis</i>	P	P	P	P	P	P	P	P		P	
<i>Lactuca saligna</i> *		P			P	P					
<i>Lactuca serriola</i> *	P	P			P	P			P	P	
<i>Lagurus ovatus</i> *			P			P		P		P	
<i>Lamprothamnium papulosum</i>					P				P	P	
<i>Lemna</i> sp.	P			P				P			
<i>Lobelia anceps</i>										P	
<i>Lolium</i> sp.*	P	P	P	P	P	P	P	P	P	P	P
<i>Ludwigia peploides</i>	P	P									
<i>Lycium ferocissimum</i> ***						P		P	P	P	
<i>Lycopus australis</i>		P									
<i>Malva parviflora</i> *						P			P		
<i>Marrubium vulgare</i> **					P						
<i>Medicago</i> sp.*		P		P	P	P	P	P	P	P	P
<i>Melaleuca halmaturorum</i>		P								P	
<i>Melilotus indica</i> *		P			P			P	P		
<i>Mentha</i> spp.*		P									
<i>Mimulus repens</i>		P		P		P	P	P	P	P	
<i>Duma florulenta</i>		P				P				P	

Species	Angas and Bremer Mouths	Dunns Lagoon	Goolwa Channel Drive	Hunters Creek	Loveday Bay	Milang	Narrung	Point Sturt	Poltalloch	Teringie	Waltowa
<i>Myriophyllum caput-medusae</i>		P		P							
<i>Myriophyllum salsugineum</i>	P	P		P	P	P					
<i>Oxalis pes-caprae</i> **						P				P	
<i>Paspalum distichum</i> *	P	P	P	P	P	P		P	P	P	
<i>Pennisetum clandestinum</i> *	P										
<i>Persicaria lapathifolia</i>	P	P									
<i>Phragmites australis</i>	P	P	P			P	P				
<i>Phyla canescens</i> *					P						
<i>Picris angustifolia</i> ssp. <i>angustifolia</i>		P			P						
<i>Plantago coronopus</i> *	P	P	P	P	P	P		P		P	
<i>Poa annua</i> *											P
<i>Polygonum aviculare</i> *								P			
<i>Polypogon monspeliensis</i> *	P		P	P	P	P	P	P	P	P	
<i>Potamogeton crispus</i>		P		P							
<i>Potamogeton pectinatus</i>		P	P	P			P				
<i>Puccinellia</i> spp.*							P				
<i>Ranunculus trichophyllus</i> *					P						
<i>Ranunculus trilobus</i> *					P						
<i>Reichardia tingitana</i> *	P	P	P		P	P		P		P	P
<i>Rorippa nasturtium-aquaticum</i> *	P										
<i>Rorippa palustris</i> *							P				
<i>Rumex bidens</i>	P	P			P	P					
<i>Ruppia megacarpa</i>				P							
<i>Ruppia polycarpa</i>			P	P		P	P	P	P	P	
<i>Ruppia tuberosa</i>				P	P		P		P	P	
<i>Salsola australis</i>										P	
<i>Samolus repens</i>		P	P		P	P	P	P		P	

Species	Angas and Bremer Mouths	Dunns Lagoon	Goolwa Channel Drive	Hunters Creek	Loveday Bay	Milang	Narrung	Point Sturt	Poltalloch	Teringie	Waltowa
<i>Sarcocornia quinqueflora</i>		P	P	P	P	P	P	P	P	P	P
<i>Scabiosa atropurpurea</i> *		P				P					
<i>Schoenoplectus pungens</i>		P	P	P	P						
<i>Schoenoplectus validus</i>		P		P	P						
<i>Senecio pterophorus</i> *	P	P	P	P	P	P	P	P		P	P
<i>Senecio runcinifolius</i>		P									P
<i>Senecio</i> sp.	P					P	P			P	P
<i>Silybum marianum</i> **					P	P					
<i>Sonchus asper</i> *		P				P					
<i>Sonchus oleraceus</i> *	P	P	P	P	P	P	P	P	P	P	P
<i>Spergularia brevifolia</i> *		P	P		P	P	P	P	P	P	P
<i>Suaeda australis</i> *		P	P	P	P	P	P	P	P	P	P
<i>Tecticornia pergranulata</i> ssp. <i>pergranulata</i>											P
<i>Trifolium</i> sp.*	P	P			P	P		P		P	
<i>Triglochin procerum</i>	P	P									
<i>Triglochin striatum</i>	P	P	P	P		P	P	P		P	
<i>Typha domingensis</i>	P	P	P	P	P	P					
<i>Urtica urens</i> *		P									
<i>Vallisneria australis</i>	P	P		P							
<i>Vicia sativa</i> *						P					
<i>Wilsonia rotundifolia</i>		P	P		P	P					
Species Total	36	72	32	38	53	64	32	40	26	51	18
Exotics Total	25	38	18	16	32	41	14	26	16	27	11
% Exotics	69.44	52.78	56.25	42.11	60.38	64.06	43.75	65	61.53	52.94	61.11

Appendix 5: Taxa present (P) at lakeshore sites from spring 2008 to autumn 2013 (*denotes exotic taxon; **denotes proclaimed pest plant in South Australia; ***denotes weed of national significance; #denotes listed as rare in South Australia)

Species	Lake Albert	Lake Alexandrina	Goolwa Channel
<i>Acacia myrtifolia</i>	P		
<i>Anagallis arvensis</i> *	P		
<i>Apium graveolens</i> *		P	
<i>Arctotheca calendula</i> *	P	P	
<i>Asparagus officinalis</i> *			P
<i>Aster subulatus</i> *	P	P	P
<i>Atriplex prostrata</i> *		P	P
<i>Atriplex</i> spp.		P	P
<i>Atriplex suberecta</i>		P	
<i>Avena</i> spp.*	P	P	
<i>Azolla filiculoides</i>		P	P
<i>Berula erecta</i> *		P	P
<i>Bolboschoenus caldwellii</i>	P	P	P
<i>Brassica rapa</i> *		P	
<i>Brassica tournifortii</i> *		P	P
<i>Briza minor</i> *		P	
<i>Bromus catharticus</i> *	P		
<i>Bromus diandrus</i> *	P	P	P
<i>Bromus hordeaceus</i> *	P	P	P
<i>Calystegia sepium</i>		P	P
<i>Carex fascicularis</i>		P	
<i>Centaurea calcitrapa</i> *	P	P	P
<i>Centaurium tenuiflorum</i> *		P	
<i>Ceratophyllum demersum</i> #		P	P
<i>Chara</i> spp.		P	
<i>Chenopodium album</i> *	P		
<i>Chenopodium glaucum</i> *	P	P	P
<i>Chenopodium nitariaceum</i>		P	
<i>Conyza bonariensis</i> *	P	P	P
<i>Cotula coronopifolia</i> *	P	P	P
<i>Cyperus exaltatus</i>			P
<i>Cyperus gymnocaulos</i>	P	P	P
<i>Distichlis distichophylla</i>	P	P	P
<i>Duma florulenta</i>	P	P	P
<i>Ehrharta longiflora</i> *	P	P	
<i>Einadia nutans</i>		P	
<i>Eleocharis acuta</i>		P	P
<i>Enchylaena tomentosa</i>	P	P	P
<i>Epilobium pallidiflorum</i>	P		P
<i>Eragrostis australasica</i>	P		
<i>Eragrostis curvula</i> **	P	P	
<i>Eragrostis</i> sp.		P	P

Species	Lake Albert	Lake Alexandrina	Goolwa Channel
<i>Euphorbia terracina</i> *	P		
<i>Ficinia nodosa</i>	P	P	P
<i>Foeniculum vulgare</i> *		P	
<i>Fumaria bastardii</i> *		P	
<i>Glyceria australis</i>		P	
<i>Helichrysum luteoalbum</i>		P	
<i>Holcus lanatus</i> *		P	
<i>Hordeum vulgare</i> *	P	P	
<i>Hydrocotyle verticillata</i>		P	P
<i>Hypochaeris glabra</i> *	P	P	
<i>Hypochaeris radicata</i> *	P	P	
<i>Isolepis platycarpa</i>	P	P	
<i>Isolepis producta</i>	P	P	
<i>Isolepis</i> sp.	P	P	
<i>Juncus acutus</i> *		P	
<i>Juncus kraussii</i>		P	P
<i>Juncus subsecundus</i>		P	P
<i>Juncus usitatus</i>		P	P
<i>Lachnagrostis filiformis</i>	P	P	P
<i>Lactuca saligna</i> *		P	P
<i>Lactuca serriola</i> *	P	P	
<i>Lagurus ovatus</i> *	P	P	
<i>Lemna</i> sp.		P	P
<i>Limosella australis</i>		P	
<i>Lobelia anceps</i>		P	
<i>Lolium</i> spp.*	P	P	P
<i>Ludwigia peploides</i>		P	
<i>Lupinus cosentinii</i> *			P
<i>Lycopus australis</i>		P	P
<i>Lythrum salicaria</i>			P
<i>Medicago</i> spp.*	P	P	P
<i>Melaleuca halmaturorum</i>	P		
<i>Melilotus indica</i> *	P	P	P
<i>Mentha australis</i>		P	P
<i>Mentha</i> spp.*		P	P
<i>Mimulus repens</i>	P		P
<i>Myriophyllum salsugineum</i>		P	P
<i>Onopordum acanthium</i> *		P	
<i>Paspalum distichum</i> *	P	P	P
<i>Pennisetum clandestinum</i> *	P	P	P
<i>Persicaria lapathifolia</i>		P	
<i>Phragmites australis</i>	P	P	P
<i>Picris angustifolia</i> ssp. <i>angustifolia</i>		P	P
<i>Plantago coronopus</i> *	P	P	P
<i>Plantago lanceolata</i> *			P

Species	Lake Albert	Lake Alexandrina	Goolwa Channel
<i>Plantago major</i> *			P
<i>Polygonum aviculare</i> *		P	P
<i>Polypogon monspeliensis</i> *	P	P	P
<i>Potamogeton crispus</i>			P
<i>Potamogeton pectinatus</i>	P	P	P
<i>Puccinellia</i> spp.	P	P	
<i>Ranunculus trilobus</i> *		P	P
<i>Reichardia tingitana</i> *	P	P	
<i>Rorippa nasturtium-aquaticum</i> *		P	
<i>Rorippa palustris</i> *		P	
<i>Rumex bidens</i>	P	P	P
<i>Ruppia tuberosa</i>		P	
<i>Salix babylonica</i> *		P	P
<i>Samolus repens</i>			P
<i>Sarcocornia quinqueflora</i>	P	P	
<i>Scabiosa atropurpurea</i> *			P
<i>Scaevola</i> sp.	P		
<i>Schoenoplectus pungens</i>	P	P	P
<i>Schoenoplectus validus</i>	P	P	P
<i>Sclerolaena blackiana</i>		P	
<i>Senecio cunninghamii</i>		P	
<i>Senecio pterophorus</i> *	P	P	
<i>Senecio runcinifolius</i>		P	
<i>Senecio</i> sp.		P	P
<i>Silybum marianum</i> **		P	P
<i>Solanum nigrum</i> *		P	P
<i>Solanum</i> spp.		P	
<i>Sonchus asper</i> *		P	
<i>Sonchus oleraceus</i> *	P	P	P
<i>Spergularia brevifolia</i> *	P	P	
<i>Suaeda australis</i> *	P	P	P
<i>Tamarix aphylla</i> ***		P	
<i>Trifolium</i> sp.*	P	P	
<i>Triglochin procerum</i>		P	P
<i>Triglochin striatum</i>	P	P	P
<i>Triticum</i> spp.*		P	
<i>Typha domingensis</i>		P	P
<i>Urtica urens</i> *		P	
<i>Vallisneria australis</i>			P
<i>Vicia sativa</i> *	P	P	
<i>Wilsonia rotundifolia</i>		P	
Species total	58	108	70
Exotic Total	34	54	30
% Exotics	58.62	49.54	42.86