

# District Weed Management Guide Port Augusta - Quorn



November 2023

# Disclaimer

The South Australian Arid Lands Landscape Board, and its employees do not warrant or make any representation regarding the use, or results of use of the information contained herein as to its correctness, accuracy, reliability, currency or otherwise. The South Australian Arid Lands Landscape Board and its employees expressly disclaim all liability or responsibility to any person using the information or advice.

This work is copyright. Apart from any use permitted under the Copyright Act 1968 (Commonwealth), no part may be reproduced by any process without prior written permission obtained from the South Australian Arid Lands Landscape Board. Requests and enquiries concerning reproduction and rights should be directed to the General Manager, South Australian Arid Lands Landscape Board, Old Court House, PO Box 297, Port Augusta, SA, 5700

© South Australian Arid Lands Landscape Board 2022


Photo front page: African boxthorn (Credit: P Hodges), Devil's rope cactus (Credit: P Hodges) and African rue (Credit: DWLBC).

# Purpose of the Guide

This guide provides information on priority weeds in the Port Augusta - Quorn District within the SA Arid Lands (SAAL) Landscape Board region. The intent is to guide prioritisation of weed management activities within the district over the next five years.

This document outlines management actions that landholders and organisations can implement within the District with the aim to reduce the current and potential impacts of priority weeds. These management actions, outlined for each of the priority weeds, are in accordance with South Australian State policies for declared weeds and the SAAL Landscape Board regional weed policies.

It also provides information regarding the known locations, possible threats, impacts, policy and monitoring activities for each of the priority weeds and provides a link to current best practice control methods.



Weed species included in this guide are at differing stages on the invasion curve. All of the included weeds have a demonstrated ability to rapidly expand their distribution given favourable seasonal conditions. Preventing the establishment of new infestations and eradication of localised populations is considered the most efficient use of resources. Conversely, trying to control widespread, well-established, persistent weeds can quickly exhaust resources. Keeping this in mind we have sought to prioritise weeds that can be controlled before they become too widespread to contain.

The District groups can be influential in the implementation of on-ground land management programs within their district. By following this guide, landholders, group members and organisations can develop proactive weed management programs within their district for the long term protection of its environmental and productive assets.

# Role of the Landscape Board

The SA Arid Lands Landscape Board have a strong commitment to weed control and have a responsibility under the Landscape SA Act 2019 to ensure this is carried out by landholders across the region.

Weed species that are declared under the Landscape Act requiring landholders to control on their land can be found at [https://www.pir.sa.gov.au/biosecurity/weeds/landholder\\_responsibilities](https://www.pir.sa.gov.au/biosecurity/weeds/landholder_responsibilities).

The Board staff are available to provide advice and assistance on weed issues and coordinate projects addressing identified priority weed control when funding is available.

## Roadsides Management

Road reserves are a recognised pathway for the introduction and movement of declared plants. It is a priority of the SA Arid Lands Landscape Board to stop the establishment and minimise the spread of weeds by managing road reserves through a systematic inspection and strategic control regime.

The Landscape Board is responsible for the control of declared weeds on public roadsides outside of townships and may recover costs from the adjacent landholder. Further information on management of roadsides can be found at [https://www.pir.sa.gov.au/\\_data/assets/pdf\\_file/0009/388593/manual-for-roadside-weed-management.pdf](https://www.pir.sa.gov.au/_data/assets/pdf_file/0009/388593/manual-for-roadside-weed-management.pdf). Weed species that are undeclared on roadsides are not the responsibility of the SA Arid Lands Landscape Board. Further information can be obtained from the local council.

Limited resources determine that not all declared plants will be managed on road reserves to the same levels. A plant's level of management will be determined by the Weed Risk Assessment system and the potential threat that it poses to the adjoining land-use.

The following plants have been identified as priorities for roadside control in the Port Augusta-Quorn District:

- Buffel grass
- African boxthorn
- African rue

## New Infestations or Unknown Weeds

If you find a new infestation of a weed you haven't had before or a weed species that you don't know, the Landscape Board would like to know about it. You can contact the SA Arid Lands Landscape Board on (08) 8429 9666 and ask for the Community Landscape Officer for your District or the Biosecurity Officer.

If you could provide the following information when you call, this will help us assist you:

- Your name
- Name of property (if applicable)
- Paddock name (if applicable)
- GPS location or description of location
- Description of weed
- Extent of weed
- Description of site where weed is located

It would also help if you could take some photos of the weed. The Officer you speak to will be able to provide an email address for the photos to be sent to. We will then attempt to identify the weed and we may forward the information and photo to the Weeds Botanist at the State Herbarium.

# Port Augusta - Quorn District

The Port Augusta – Quorn (PAQ) district is a recent addition to the SA Arid Lands Landscape region, as of July 2020, and includes diverse landscapes and communities. The new district stretches from Cultana on the Eyre Peninsula to Port Augusta, and follows the range from Mt Brown, north, through the Arden Vale ranges.

Geographically the area of 538,000 hectares is characterised by parallel ranges and gorges, a mixture of plains and tablelands at the head of the Spencer Gulf, and 122 kilometres of coastline and coastal habitat. The diverse landscape supports productive agricultural grazing and marginal cropping, rich biodiverse habitats and vibrant towns with a total population of 15,000 residents in the district.

## Declared Weeds

The weeds identified as priorities for the Port Augusta - Quorn District dealt with in this guide are declared weeds under the *Landscape SA Act 2019* and demonstrated to be regional threats, where their management is a strategic option.

Weed risk assessments have been undertaken by the SAAL Landscape Board on each of the priority weeds. This process determines a weed's risk as low, medium or high, in a particular land system, by assessing the weeds invasiveness, impact and potential distribution. The management guide for each weed was identified using the assessment of each weed's risk category in the table on the following page.

## Alert Weeds

These declared weeds have not been sighted in the Port Augusta - Quorn District. However, there have been infestations of them in neighbouring district(s) and the threats they pose are significant. If you discover any of these plants in the Port Augusta - Quorn District please contact SA Arid Lands Landscape Board on (08) 8429 9666 immediately and report the location. The following links are provided for information and identification:

- [Noogoora burr \(\*Xanthium strumarium\*\)](#)

## Established Weed Species

Weed species that are widespread and/or established, will be managed if opportunity arises through project funding or on a case by case situation, with the same objective as this plan. Landholders are encouraged to contain any weed on their land to prevent further spread. The following links are provided for information, identification and/or policies:

- [Bathurst burr \(\*Xanthium spinosum\*\) – Declared Plant Policy](#)
- Castor oil plant (*Ricinus communis*) – Not declared – [Plant Policy](#)
- Mexican poppy (*Argemone ochroleuca*) – Not declared ([See Appendix 1](#))
- Pepper tree (*Schinus molle*) – Not declared ([See Appendix 1](#))
- [Salvation Jane \(\*Echium plantagineum\*\) – Declared Plant Policy](#)

# Priority weeds in the Port Augusta - Quorn District

Priority Weed	Management guide	Distribution in the District	Description of management guide
African boxthorn ( <i>Lycium ferocissimum</i> )	Manage weed	Common	Reduce overall economic and/or social impacts through targeted management. <a href="#">See page 9 for more details.</a>
African rue ( <i>Peganum harmala</i> )	Protect sites	Limited	Prevent spread of the weed species to key sites/assets of high economic, environmental and/or social value. <a href="#">See page 12 for more details.</a>
Buffel grass ( <i>Cenchrus ciliaris</i> )	Protect sites	Restricted to certain areas	Prevent spread of the weed species to key sites/assets of high economic, environmental and/or social value. <a href="#">See page 15 for more details.</a>
Cactus species ( <i>Austrocylindropuntia</i> , <i>Opuntia</i> and <i>Cylindropuntia</i> <i>spp.</i> )	Destroy infestations	Common	Aims to significantly reduce the extent of the species. <a href="#">See page 18 for more details.</a>
Innocent weed ( <i>Cenchrus longispinus</i> )	Protect Sites	Limited	Prevent spread of the weed species to key sites/assets of high economic, environmental and/or social value. <a href="#">See page 25 for more details.</a>
Khaki weed ( <i>Alternanthera pungens</i> )	Destroy Infestations	Limited	Significantly reduce the extent of the species. <a href="#">See page 28 for more details.</a>
Mesquite ( <i>Prosopis sp.</i> )	Destroy infestations	Very limited	Significantly reduce the extent of the species. <a href="#">See page 31 for more details.</a>
Parkinsonia ( <i>Parkinsonia aculeata</i> )	Destroy infestations	Very limited	Significantly reduce the extent of the species. <a href="#">See page 35 for more details.</a>

# Definitions of management guide aims

---

<b>Management guide</b>	<b>Aims</b>
<b>Alert</b>	<p>Species that are not known to be present in the management area and present a significant threat if permitted to enter and establish.</p> <p><b>Aims to prevent the species arriving and establishing:</b></p> <ul style="list-style-type: none"><li>• Prevention of entry to management area</li><li>• Ongoing surveillance for incursions</li><li>• Training &amp; awareness activities for the community to enable early detection</li><li>• Report and destroy any plants found</li></ul>
<b>Eradicate</b>	<p><b>Aims to remove the species:</b></p> <ul style="list-style-type: none"><li>• Detailed surveillance &amp; mapping to locate all infestations</li><li>• Destruction of all infestations including seed banks</li><li>• Prevention of entry to management area and movement within</li><li>• Must not be grown and all cultivated plants to be removed</li><li>• Monitor progress towards eradication</li></ul>
<b>Destroy</b>	<p><b>Aims to significantly reduce the extent of the species:</b></p> <ul style="list-style-type: none"><li>• Detailed surveillance &amp; mapping to locate all infestations</li><li>• Destruction of all infestations, aiming for local eradication at feasible sites.</li><li>• Prevention of entry to management area and movement and sale within</li><li>• Must not be grown</li><li>• Monitor progress towards reduction</li></ul>
<b>Contain spread</b>	<p><b>Aims to prevent the ongoing spread of the species:</b></p> <ul style="list-style-type: none"><li>• Surveillance &amp; mapping to locate all infested properties</li><li>• Control all infestations, aiming for a significant reduction in weed density</li><li>• Prevention of entry to management area and movement and sale within</li><li>• Must not allow to spread from cultivated plants (if grown)</li><li>• Monitor change in current distribution</li></ul>
<b>Protect sites</b>	<p><b>Aims to prevent spread of the weed species to key sites/assets of high economic, environmental and/or social value:</b></p> <ul style="list-style-type: none"><li>• Weed may be of limited current distribution but only threatens limited industries/habitats (lower weed risk). Or the weed may be more widespread but is yet to invade/impact upon many key industries/habitats (higher weed risk)</li><li>• Surveillance &amp; mapping to locate all infested properties</li><li>• Identification of key sites/assets in the management area</li><li>• Control of infestations in close proximity to key sites/assets, aiming for a significant reduction in weed density.</li><li>• Limits on movement within the management area</li><li>• Must not allow to spread from cultivated plants (if grown) in close proximity to key sites/assets</li><li>• Monitor change in current distribution within and in close proximity to key sites/assets</li></ul>
<b>Manage weed</b>	<p><b>Aims to reduce the overall economic, environmental and/or social impacts of the weed species through targeted management</b></p>

- Research and develop integrated weed management (IWM) packages for the species, including herbicides and biological control, where feasible
- Promote IWM packages to landholders
- Monitor decrease in weed impacts with improved management
- Identify key sites/assets in the management area and ensure adequate resourcing to manage the weed species

**Manage sites**

**Aims to maintain the overall economic, environmental and/or social value of key sites/assets through improved general weed management.**

- Promote general IWM principles to landholders, including the range of control techniques, maintaining competitive vegetation/crops/pastures, hygiene & property management plans
- Identify key sites/assets in the management area & ensure adequate resourcing to manage these to maintain their values
- Broaden focus beyond weeds to all threatening processes

**Monitor**

**Aims to detect any significant changes in the species' weed risk**

- Monitor the spread of the species and review any perceived changes in weediness

**Limited action**

**The weed species would only be targeted for coordinated control in the management area if its local presence makes it likely to spread to land uses where it ranks as a higher priority**

- Undertake control measures if required for the benefit of other land uses at risk
- Otherwise limited advice to land managers, if required



# African boxthorn *Lycium ferocissimum*

## Description

Branched shrub to 5 m high and 3 m wide. Spines occur on the main stems and branchlets, and branchlets terminate with a spine. Slightly fleshy leaves, 10-40 mm long, in clusters at the nodes. Flowers are pale lilac to white with purple markings at the base. Flowers predominately in summer but may occur all year round. Fruit is dull orange-red berry, 1 cm diameter. Originally from South Africa, African boxthorn was introduced in 1845 as a hedge plant. The fruit is a breeding place for numerous insects including common house fly, fruit fly and dried fruit beetles.

**Do not confuse** African boxthorn (*Lycium ferocissimum*) with the native [Australian boxthorn \(\*Lycium australe\*\)](#).



Leaves and spines (Credit: J Pitt)



Immature and mature fruit (Credit: J Pitt)

## Distribution

**Australian distribution:** All states

**SA distribution:** Common in large areas of the arid lands and along and near the coast of Eyre Peninsula.

**District distribution:** Common at low densities throughout the district.

**Potential distribution:** Has the potential to infest watercourses and flood plains, especially when boxthorns are located upstream. Boxthorns may increase in density where unmanaged.

## Threats and Impacts

### Invasiveness

African boxthorn can only be dispersed through seed, which is eaten by birds and foxes and viable when excreted. Seed can be easily dispersed over large areas, particularly under trees, poles and fences where birds have perched. Regrowth may occur from broken roots and cut stumps following physical control.

### Impacts

African boxthorn may form dense stands which compete with native species and pastures. The sharp spines prevent grazing and make stock movement difficult, particularly near watercourses which may prevent access.

African boxthorn can impact on native fauna by reducing habitat quality. Thickets of African boxthorn can harbour feral pests such as foxes and rabbits.

### Persistence

African boxthorn has the ability to reshoot and reinvade areas where control has been undertaken, therefore long-term follow-up is required. Reinfestation of boxthorn may occur from untreated plants in surrounding areas.

### Policy

#### National Strategy

African boxthorn is classified as a Weed of National Significance (WONS). The extent and potential extent in Australia will be well understood, stakeholders well equipped with information tools and knowledge, and strategic management will contain and reduce the weed's impacts.

#### State Policy

To protect native vegetation and maintain access to pasture throughout SA, under the *Landscape SA Act 2019*:

- Restrict movement of African boxthorn on public roads
- Prohibit sale of African boxthorn or produce or goods carrying African boxthorn
- Require landholders to destroy African boxthorn and inhibit its propagation as far as is reasonably achievable
- Allow Landscape Boards to recover costs of control of African boxthorn on road reserves from adjoining landholders.

#### SA Arid Lands Landscape Board Risk Assessment

The SAAL Landscape Board risk assessment for African boxthorn is to **MANAGE WEED**.

Aim: To reduce the overall economic, environmental and/or social impacts of the weed species through targeted management.

#### Port Augusta - Quorn District Risk Management Action

As above.

### Best Practice Control

Control method	Best time to control	Link for PIRSA current best practice chemical control and rates
<b>Foliar Spray</b>		
Useful for all sized plants. The entire plant is sprayed when actively growing, i.e. when the plant has green healthy leaves. Plants should not be sprayed when they are stressed or when plants do not have foliage. Boxthorns are deciduous plants and will shed their leaves during winter and/or when stressed.	Summer and Spring when actively growing	<a href="#">African boxthorn - PIRSA</a>
<b>Basal Bark</b>		
Useful for all sized plants Spray all sides of each stem with the recommended herbicide mix, to a height	Summer and Spring when actively growing	<a href="#">African boxthorn - PIRSA</a>

---

of between 250 – 750 mm above soil level.

---

### **Cut Stump**

---

Useful for all sized plants

Cut the entire plant down at the base using a chainsaw and immediately (within 10 seconds) spray or paint the recommended herbicide mix onto the cut stump.

Summer and Spring when actively growing

[African boxthorn - PIRSA](#)

---

### **Physical and/or Organic Control**

---

Small plants can be hand-pulled or dug up. Large plants can be removed using heavy machinery.

Best done when there is ground moisture as the plants will be more readily removed.

Uprooted plants should be piled & destroyed by burning as they can provide shelter for vermin and the spines can still cause injury.

After removal, the area should be monitored for regrowth from root fragments left in the soil, and for germinating seedlings.

---

## **Recommended Actions for African boxthorn Management in Port Augusta - Quorn District**

1. Support landholder education in identification, monitoring and control of African boxthorn.
2. Encourage landholders to provide data on distribution so that it can be surveyed and mapped.
3. Provide input into identification of key sites for control of African boxthorn infestations.
4. Assist in identifying and coordinating African boxthorn control programs using integrated weed management, especially in areas where key sites are threatened.
5. Encourage landholders to monitor success, following control of African boxthorn, and carry out follow up control as necessary.
6. Encourage landholders to use best practice hygiene, to prevent movement and spread of African boxthorn.

### **Surveys and Monitoring**

Keep an eye out for African boxthorn during routine pastoral management activities. Best time for monitoring is during spring and summer as well as in the weeks and months after rainfall events when the leaves are bright green.

# African rue *Peganum harmala*

## Description

Perennial herb or shrubby plant 30-80 cm high. Leaves 1-5 cm long, bright green, divided several times into three or more linear segments. Flowers with five white broad petals, 12-17 mm long, in late spring to early summer. Fruit is slightly flattened capsule, 8-12 mm across and 7-12 mm long, which opens at the top, containing black angular seeds. Originally from the Mediterranean region and the Middle East, introduced in the 1930s for unknown reasons. It is considered to be an aphrodisiac in India, and its seeds and leaves have been used traditionally to treat various ailments including asthma, jaundice, colic and as a diuretic.



African rue plant (Credit: B Shepherd)



African rue flowers (Credit: DWLBC)

## Distribution

**Australian distribution:** Confined to New South Wales and South Australia.

**SA distribution:** Known to occur in patches at Tintinara, Taylorville, Snowtown and the mid-north as well as numerous other sites. In the SAAL region African rue primarily occurs along roadsides and flood out areas. A core infestation exists in the eastern pastoral district and other occurrences exist in the northern Flinders Ranges and in the area between Lake Torrens and Port Augusta including along the Stuart Highway.

**District distribution:** Mostly on creek-lines in the Willochra catchment. Scattered distribution identified during January 2021 roadside weed surveys at creek crossings around Quorn.

**Potential distribution:** African rue has the potential to become widespread throughout the SAAL region in disturbed areas such as roadsides and areas receiving run-on water such as flood outs and depressions. The zone of potential distribution has been described as between the 350 mm and 175 mm isohyets.

## Threats and Impacts

### Invasiveness

Dispersal of the plant is predominately through seed, with the majority being dropped close to the parent plant. Seeds can easily be dispersed through water flow, but also in mud moved by animals or vehicles. Stock and other animals may also eat the fruit and aid in the dispersal. African rue prefers disturbed sites with little or no competition, and requires moisture for seed germination. African rue is drought and salt tolerant, and therefore has the potential to thrive in semi-arid and arid areas.

### Impacts

African rue can be toxic containing more than 25 alkaloids, however the plant is highly unpalatable to livestock with few poisonings reported. The likely impact on native vegetation is unknown, but due to its tolerance to drought and salt the potential impacts could be significant.

### Persistence

African rue is difficult to destroy once established as regeneration may occur from severed root pieces. African rue is known to persist despite treatment with herbicide or manual removal.

## Policy

### National Strategy

African rue is not classified as a Weed of National Significance, therefore there is no National Strategy for this species.

### State Policy

African rue is a declared species under the *Landscape SA Act 2019*, to prevent establishment in pastoral lands:

- Prohibit movement of African rue on roads
- Prohibit the sale of African rue or contaminated produce
- Require landholders to **notify** Landscape Board of African rue infestations
- Require landholders to control African rue on their properties
- Allow recovery of roadside control costs of African rue from adjoining landholders

### SA Arid Lands Landscape Board Risk Assessment

The SAAL Landscape Board risk assessment for African rue is to **PROTECT SITES**.

Aim: To prevent spread of African rue to key sites/assets of high economic, environmental and/or social value.

### Port Augusta - Quorn District Risk Management Action

As above

### Best Practice Control

Control method and description	Best time to control	Link for PIRSA current best practice chemical control and rates
<b>Foliar Spray</b>		
Useful on all sized plants. African rue has very deep roots and glyphosate may not effectively translocate to kill established plants. Repeated application over successive seasons is required. Spray the entire plant until run off. Better results will occur on plants that have dust free foliage. Spot spraying with a knap sack or quad/ute mounted spray unit can be used for scattered occurrences. Boom spray can be used for dense infestations where no native vegetation exists between African rue plants e.g. along roads.	After rainfall events and retreat annually. Best sprayed before the plant flowers.	<a href="#">African rue - PIRSA</a>
<b>Cut stump</b>		
Cut off close to the ground and apply herbicide to stump surface immediately.	When actively growing.	<a href="#">African rue - PIRSA</a>
<b>Physical and/or Organic Control</b>		
Due to its deep roots, African rue cannot be successfully controlled through hand-pulling or grubbing (it will regrow from the broken root). Organic properties - To provide long term suppression and reduce African rue proliferation, increasing competition from native plants, by reducing grazing pressure from livestock, rabbits and kangaroos, in areas where African rue is present, is the best option.		

## **Recommended Actions for African rue Management in Port Augusta - Quorn District**

- 1.** Support landholder education in identification, monitoring and control of African rue.
- 2.** Encourage landholders to provide data on distribution of African rue so that it can be surveyed and mapped.
- 3.** Provide input into identification of key sites for control of African rue infestations.
- 4.** Assist in identifying and coordinating African rue control programs using integrated weed management, especially in areas where key sites are threatened.
- 5.** Encourage landholders to monitor success, following control of African rue, and carry out follow up control as necessary.
- 6.** Restrict sale and movement of African rue within SAAL Landscape region.

### **Surveys and Monitoring**

Undertake targeted surveys and/or opportunistic monitoring of African rue during pastoral management activities, especially along and adjacent to roadsides and other disturbed areas. Targeted surveys and control activities should occur annually, especially 2-3 weeks after rainfall. Record locations with GPS or appropriate app and advise SA Arid Lands Landscape Board of infestation.

# Buffel grass *Cenchrus ciliaris*

## Description

Buffel grass is a perennial grass to approximately 1 m high, with very strong root stock. Leaves bluish-green, 3-25 cm long, 1-6 mm wide. Leaf blade base with a row of short hairs, 0.2-2 mm long. Flower heads form dense hairy cylindrical spikes 2-15 cm long, pale or purplish. Flowers in summer. Originally from Africa and south western Asia and introduced in the 1840s as a pasture species and for erosion control. It can withstand heavy grazing and is the most drought tolerant introduced grass species in Australia.

Buffel grass may be confused with other grass species. For more information about telling the difference please click [here](#).



Buffel grass seed head (Credit: B Shepherd)



Buffel grass infestation (Credit: B Shepherd)



Buffel grass plant (Credit: B Shepherd)

## Distribution

**Australian distribution:** Common throughout central Australia. Widely cultivated by pastoralists as a preferred pasture species for cattle. In some outback towns, namely Alice Springs it was cultivated for dust control and has since successfully naturalised and overrun large tracts of land.

**SA distribution:** Large areas in the far north west of northern SA are infested with Buffel grass and it is common along the Stuart Highway (from the Northern Territory border down to Glendambo), the Tarcoola Road, Coober Pedy, Port Augusta, the Augusta Highway from Port Augusta to Mambray Creek (including rail corridor) and the rail corridor on the Port Augusta-Whyalla line. Buffel grass also occurs along selected intermittent watercourses of the SAAL region with infestations ranging from scattered single occurrences to dense patches. It also occurs in other areas including around townships and as isolated patches along secondary roads.

**District distribution:** Augusta highway from Port Augusta to Mambray Creek (including rail corridor), Port Augusta township and rural blocks, rail corridor on the Port Augusta-Whyalla line and scattered roadsides in the district.

**Potential distribution:** Buffel grass establishes readily and has the capacity to expand across a large proportion of the SA Arid Lands and neighbouring districts if left unchecked.

## Threats and Impacts

### Invasiveness

Buffel grass is easily distributed by wind, water, stock and machinery. High levels of disturbance, such as flood, fire and heavy grazing, can assist the establishment. Buffel grass can also root from lower nodes. The species requires summer rain for growth, and is not frost tolerant.

### Impacts

Buffel grass competes with and displaces native species. It forms monospecific stands, out-competing native grasses. Plants are fire resistant but can carry fire in areas where fire is not normally part of the ecosystem. Buffel grass aids fire by increasing the intensity and frequency of natural fire regimes to the point of removing competing shrubs and trees.

### Persistence

Buffel grass is the most drought tolerant introduced grass species in Australia, is highly persistent on lightly textured soils and is quick to respond to small amounts of rainfall.

## Policy

### National Strategy

Buffel grass is not classified as a Weed of National Significance, therefore there is no National Strategy for this species.

### State Policy

Buffel grass is a declared species under the *Landscape SA Act 2019*. Buffel grass must be contained to minimise its impacts on native vegetation, grazing systems, remote communities and infrastructure.

Objectives:

- Prohibit entry to area
- Prohibit movement of Buffel grass on public roads
- Prohibit the sale of Buffel grass or contaminated produce
- Landowners to control Buffel grass on their properties
- Recovery of control costs on adjoining road reserves by the Landscape Board

### SA Arid Lands Landscape Board Risk Assessment

All SA Arid Lands Landscape Districts (except Marla-Oodnadatta) lie within Zone 2 (**PROTECT SITES**) in the South Australia Buffel grass Strategic Plan 2019-2024. The Marla-Oodnadatta Landscape District lies within Zone 1 (**MANAGE WEED**) in the South Australia Buffel grass Strategic Plan 2019-2024.

Aim: Zone 1 – To reduce the overall economic, environmental and/or social impacts of Buffel grass through targeted management. Zone 2 - To prevent spread of Buffel grass to key sites/assets of high economic, environmental and/or social value.

### Port Augusta - Quorn District Risk Management Action

Port Augusta - Quorn District management action is to **PROTECT SITES** from the spread of Buffel grass.

Aim: To prevent spread of Buffel grass to key sites/assets of high economic, environmental and/or social value.

## Best Practice Control

---

Control method and description	Best time to control	Link for PIRSA current best practice chemical control and rates
<b>Foliar Spray</b>		

---



---

Spray all sides of the entire plant.  
Ensure the entire plant is covered in the herbicide mix.

Between 2- [Buffel grass - PIRSA](#)  
4 weeks  
after warm  
season rain

Spot spraying with a knap sack or quad/ute mounted spray unit can be used for scattered occurrences.

Boom spray can be used for dense infestations where no native vegetation exists between Buffel grass plants e.g. along roads.

Buffel grass can quickly regenerate from seed and ongoing monitoring and control is required after warm season rain.

---

### **Physical and/or Organic Control**

---

Burning dried-out Buffel grass allows for better control from chemical spraying when plant re-shoots after rain (e.g. better coverage of new foliage).

Organic foliar spraying can be carried out using Bioweed (680g/L pine oil) at a rate of 20L per 100L water. Complete coverage of plant is required and should be applied when the plant is actively growing. This mixture will also kill seed it contacts both on the plant and on the ground.

Grubbing can be used for isolated occurrences or small infestations.

Dig the entire plant out of the ground.

Ongoing monitoring and follow up hand pulling/grubbing will be required annually.

---

### **Recommended Actions for Buffel grass Management in Port Augusta - Quorn District**

1. Support landholder education in identification, monitoring and control of Buffel grass.
2. Encourage landholders to provide data on distribution of Buffel grass so that it can be surveyed and mapped.
3. Provide input into identification of key sites requiring control of Buffel grass.
4. Assist in identifying and coordinating Buffel grass control programs using integrated weed management, especially in areas where key sites are threatened.
5. Encourage landholders to monitor success, following control and carry out follow up control as necessary.
6. Restrict introductions and movement of Buffel grass within SAAL Landscape region.

### **Surveys and Monitoring**

Monitor during routine pastoral management activities. Look in areas adjacent to roads, railways and beside watercourses. In the weeks following rainfall, look for a large clumpy grass with purple or black seed heads.

# Cactus species

## Coral cactus *Cylindropuntia fulgida* var. *mamillata*

### Description

Usually 1-1.5 m high and occasionally up to 3 m. Upper segments are smooth greyish to dark green, 6-70 cm long and 1-5 cm wide. Segments resemble coral as they mature. Spines (1-6) emerge from depressions, with white woolly hairs and minute bristles. Flowers dull red 2-3.5 cm wide, in late spring to summer. Fruit is yellow-green, spiny, barrel shaped. Originally from Ecuador and Peru, it is unknown when or why it was introduced. Floodwaters may damage plants and also disperse segments resulting in new infestations.



Coral cactus plant (Credit: B Shepherd)



Mature Coral cactus plant (Credit: B Shepherd)

## Devil's rope cactus *Cylindropuntia imbricata*

### Description

Usually 1-2 m high and occasionally up to 3 m, often with a woody trunk. The plant is made up of strong woody segments dark to grey-green which are rope like in appearance. Sharp spines (2-30) emerge from depressions in segments, spines, 2-3 cm long, enclosed in yellow bristles. Flowers are 3-7.5 cm wide, purple or purplish-red, in late spring to summer. The fruit is usually spineless, barrel shaped and matures to a yellow colour. Originally from southern USA and Mexico, it is unknown when it was introduced but used as an ornamental garden plant.



Mature Devil's rope cactus (Credit: J Pitt)



Devil's rope flower (Credit: J Pitt)

## Jumping cholla *Cylindropuntia prolifera*

### Description

Up to 2m high. Segments are grey-green up to 15 cm long and 5 cm wide. Segment depressions contain 6-12 spines up to 2 cm long. Flowers rose to magenta, to 4 cm wide, spring to early summer. Fruit 2-4 cm long, 2-3 cm wide, spineless occasionally in short chains. Originally from USA and Mexico, first record in NSW in 1993, unknown why introduced.



Jumping cholla plant (Credit: R Holtkamp)



Jumping cholla flower (Credit: R Holtkamp)

## Hudson pear *Cylindropuntia pallida*

### Description

Branched, 1.5 m high to 3 m wide. Segments are cylindrical up to 90 cm long and 4 cm wide. Segment depressions contain clusters of 4-8 spines, up to 3.5 cm long. Pink flowers to 5 cm wide, late spring and summer. Fruit is wider towards the tip, 2 - 4.5 cm long. Originally from Mexico and introduced in 1960s as an ornamental garden plant.



Mature Hudson pear cactus (Credit: G Patrick)



Hudson pear flower

## Engelmann's prickly pear *Opuntia engelmannii*

### Description

Up to 2m tall. Upper segments are dull mid to grey-green and oval to circular shaped, 9-26 cm long. Segment depressions contain 1-12 spines, with brown woolly hairs and short yellow-brown bristles. Yellow flowers. Fruit is pear to barrel shaped, spiny, and matures to reddish-purple. Originally from USA and Mexico, unknown when and why introduced.



Mature Engelmann's prickly pear (Credit: B Shepherd)



Pads and flower buds (Credit: B Shepherd)

## Wheel cactus *Opuntia robusta*

### Description

Usually 1-2 m tall, occasionally to 4 m generally with a well-developed trunk. Segments are circular, bluish green to bluish grey to 40 cm in diameter. Segment depressions are widely spaced containing 1-12 spines, 5 cm long with brown woolly hairs and yellow to brown bristles. Flowers are yellow ageing to white, 5-8cm diameter. Fruit is pink to purple, barrel shaped to 8 cm long and 6 cm wide. Originally from Mexico, unknown when introduced but used as an ornamental garden plant. Wheel cactus is considered rare and endangered in its native habitat and is listed on the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES).



Wheel cactus plants with flower buds (Credit: J Pitt)



Wheel cactus pads (Credit: J Pitt)

## Prickly pear *Opuntia stricta*

### Description

Usually 0.5-1m tall, and 0.5-5m across. The basal stem segments often thicken and form a trunk. The upper stem segments are dull mid to grey-green and oval shaped. Segment depressions are usually spineless or contain 1-11 spines, 1-6 cm long, with brownish woolly hairs and short yellow bristles. Flowers are yellow, 5-8 cm wide, summer. Fruit is pear shaped, smooth and purple at maturity. Originally from drier tropical and sub-tropical America, introduced before 1839 as an ornamental garden plant and for use as food for cochineal insects used to produce dye for soldiers' coats. Plants may still establish from segments following disposal of garden waste unless buried to a depth greater than 1 metre.



Prickly pear plant (Credit: P Hodges)



Prickly pear flower (Credit: P Hodges)

### Distribution

**Australian distribution:** These cacti species are common throughout Australia, both in gardens and as naturalised plants. In most states one or several cactus species are declared weeds.

**SA distribution:** There are significant infestations of Prickly pear and/or Wheel cactus near Peterborough, Terowie, along the River Murray cliffs, in the Blinman-Parachilna area of the Flinders Ranges and around Port Augusta. There is an infestation of Engelmann's prickly pear in the Flinders Ranges west of Quorn which is often confused with Wheel cactus. Devil's rope cactus occurs to a lesser extent in these areas as well as in other areas around northern SA, primarily in the mid north agricultural district, the Flinders Ranges and the Olary Ranges.

There are three sites – two east of Coober Pedy and one in the North Flinders - where Coral cactus and/or Jumping cholla have naturalised and are spreading, however these populations are small and manageable. It is likely that these are not the only occurrences in northern South Australia and that others exist possibly in abandoned gardens or around ruins. Hudson pear occurs near Port Augusta and Whyalla.

**District distribution:** There are cactus infestations in various areas in the Port Augusta - Quorn District. Prickly pear populations are throughout Port Augusta and surrounds, including Saltia Creek. Devil's rope cacti are also present in low densities in creek-lines and ranges. Engelmann's prickly pear is common through the ranges from Mt Brown to Buckaringa and also in the Arden Vale area. Wheel cacti have been found alongside the railway line in Port Augusta. Hudson pear is present west of Port Augusta on properties adjacent to the Lincoln Highway and in the township at Back Beach.

**Potential distribution:** Cacti species have the potential to establish across most of the arid and semi-arid areas of South Australia.

## Threats and Impacts

### Invasiveness

Wheel cactus, Prickly pear and Engelmann's prickly pear are highly invasive. They have high seed production, reproduce vegetatively, seeds are dispersed by birds and other animals and seedlings establish easily. Flood events have also been observed to promote spread of cacti species. Devil's rope cactus, Jumping cholla and Coral cactus are also invasive species, well adapted for vegetative spread and able to withstand drought conditions.

### Impacts

At high density, infestations of cacti species can render land unsuitable for grazing as thickets exclude livestock and can degrade biodiversity as the infestations outcompete native plants and reduce habitat for native fauna.

### Persistence

The ability of cacti to recover from control is very high. Re-establishment is assisted by the longevity of seed stored in the soil and the fact that new plants can grow from untreated or dropped pads. Follow up control is essential.

## Policy

### National Strategy

All Opuntoid cacti species *Austrocylindropuntia*, *Cylindropuntia* and *Opuntia* (except *Opuntia ficus-indica*) are classified as Weeds of National Significance. The Strategic Plan aims to deliver the following goals and objectives:

- Prevent new infestations from establishing.
- Strategically manage established infestations.
- Increase capability and commitment to manage Opuntoid cacti.

### State Policy

Prevent *Opuntia* species from competing with more desirable plants and restricting access in the pastoral areas of the State, under the *Landscape SA Act 2019*:

- Prohibit movement of *Opuntia* species on public roads and entry into SA;
- Prohibit sale of *Opuntia* species or their seeds, or contaminated material;
- Require landholders to control *Opuntia* species on their properties; and
- Allow recovery of costs of *Opuntia* species control on road reserves by Landscape Board.

### SA Arid Lands Landscape Board Risk Assessment

The SAAL Landscape Board risk assessment for *Opuntoid* and *Cylindropuntia* cacti species is to **DESTROY INFESTATIONS**.

Aim: To significantly reduce the extent of *Opuntoid* and *Cylindropuntia* cacti species in the SAAL Landscape region.

### Port Augusta - Quorn District Risk Management Action

As above

## Best Practice Control

Following initial control of mature cactus plants, cactus seeds deposited under mature plants and pads that were missed or dropped off during treatment will germinate and grow, therefore follow up monitoring and control around mature plants is required for several years.

---

### Control method and description

---

#### Biological

---

Cochineal scale (*Dactylopius* spp.) may be used as a biological control for various Opuntioidei cacti and *Cylindropuntia* species including Engelmann's prickly pear, Hudson pear, Prickly pear, Wheel cactus, Devil's rope, Coral cactus and Jumping cholla.

Evidence of the presence of cochineal appears as small white "cotton wool" spots on the cactus pads. Cochineal create these structures and live under them. They suck fluid from plant tissues and in high enough densities energy production may be impacted and the plant's growth is restricted or the plant may die.

Pads from infected plants may be removed and wedged low on a new host plant, protected from rain where possible, with the most heavily infested surface as close as possible to the surface of the new host plant. This will allow for movement of wingless cochineal to the new host plant.

**Note: Different cochineal species and lineages work on different Opuntioidei cacti species. Please see Appendix 2 for more information.**

---

Foliar spray	Best time to control	Link for PIRSA current best practice chemical control and rates
Useful for all cacti species. Suitable for larger infestations. Spray all sides of every cactus pad until chemical runs off. Spray only when cacti are green and healthy.	Any time of year providing plants are not stressed. Warmer months are preferred.	<a href="#">Opuntia cacti (prickly pear/wheel cactus) - PIRSA</a> <a href="#">Cylindropuntia cacti (devils rope, Hudson pear) - PIRSA</a>

---

Stem inject		
Useful for most cactus species. Suitable for isolated plants or small infestations. Inject a measure of herbicide into each cactus stem or in at least every 4th pad utilising a Velpar® gun and injecting lance.	Any time of year	<a href="#">Opuntia cacti (prickly pear/wheel cactus) - PIRSA</a> <a href="#">Cylindropuntia cacti (devils rope, Hudson pear) - PIRSA</a>

---

#### Physical

---

It is possible to dig out cacti but care must be taken due to their spines. This is only practical for small isolated plants. Care must be taken not to drop any segments, as these may grow and develop into new plants. Material must be disposed of via deep burial (at least 1m). This method may be used all year round. Note: It is an offence to transport Opuntioidei cacti on roads, so burial at infestation site is recommended.

#### Fire

---

Hot fires may kill plants but regrowth may occur, requiring follow-up control. Burning can assist in gaining access to large infestations allowing use of other control activities. This method may not be practical where cacti are growing amongst native vegetation.

---

## **Recommended Actions for Cactus Management in Port Augusta - Quorn District**

- 1.** Support landholder education in identification, monitoring and control of Opuntia species.
- 2.** Encourage landholders to provide data on Opuntia infestations so that it can be surveyed and mapped.
- 3.** Provide input into identification of key sites requiring control of Opuntia species.
- 4.** Assist in identifying and coordinating Opuntia control programs using integrated weed management, especially in areas where key sites are threatened.
- 5.** Encourage landholders to undertake control of all Opuntia species infestations on their property.
- 6.** SAAL to promote Cochineal nursery and trading clean cacti pads for cochineal infested pads with land managers.
- 7.** Encourage landholders to monitor success, following control of Opuntia, and carry out follow up control as necessary.
- 8.** Restrict sale and movement of all Opuntia species within SAAL Landscape region.

## **Surveys and Monitoring**

Keep an eye out for cactus plants during routine pastoral management activities. Look for old plantings of cactus in historic gardens around ruins, railway sidings, illegal dump sites, beside watercourses and areas associated with human settlement. Survey at any time of year.



# Innocent weed *Cenchrus longispinus* and Spiny burr-grass *Cenchrus incertus*

## Description

**Innocent weed** *Cenchrus longispinus* - annual grass to 70 cm. Leaves narrow 1-6 mm wide, 3-15 cm long, with a reddish or purplish tinge. Flower in summer and autumn. Seed head has 8-18 'burrs', 3-12 mm across, with 40-70 sharp spines, 3.5-7 mm long. Burrs are reddish or purplish-green when young but mature to a brown or straw colour. Burrs are on stalks 1-5 mm long. Seeds are egg shaped, 2-4 mm long and 2-3 mm wide.

**Spiny burr-grass** *Cenchrus incertus* - annual and occasionally perennial grass to 60 cm tall. Leaves narrow 2-8 mm wide to 2-20 cm long. Flower in summer and autumn. The seed-head has up to 40 'burrs', 3-10 mm across, with 8-45 sharp spines, 2-5 mm long. Burrs are purplish-green when young but mature to a brown or straw colour. Seeds are egg shaped, 2-4 mm long and 2-3 mm wide.

Originally from North and Central America, these weeds were introduced in 1886 for unknown reasons.

*Cenchrus longispinus* and *Cenchrus incertus* can be confused with each other and other *Cenchrus* sp. Species are distinguished by their 'burrs'.



Innocent weed plant (Credit: A Harvey)



Innocent weed seed heads (Credit: A Harvey)

## Distribution

**Australian distribution:** Innocent weed is widespread throughout the Murray Mallee and irrigated areas along the Murray River of both VIC and SA. It is common in coastal QLD and is found in small isolated areas of the NT and WA.

**SA distribution:** In SA, it has been found on the Eyre Peninsula, northern pastoral and agricultural districts, and in isolated patches of the South East. Major infestations are in the Riverland irrigation areas, the Northern Murray Mallee dryland areas and in urban areas.

**District distribution:** There are scattered plants of Innocent weed at various locations in Port Augusta.

**Potential distribution:** Innocent weed has the potential to establish throughout most semi-arid areas of SA.

## Threats and Impacts

### Invasiveness

Innocent weed is readily spread by dispersal of its burrs which contain the plants' seeds, as they attach to animals, vehicles and clothing. Burrs may also be distributed by water and contaminated agricultural produce, especially hay. A weed of disturbed sites such as waste areas, roadside, pastures and cultivation areas. Seedlings have been known to establish from seed deeply buried up to 25 cm in sandy soils.

### Impacts

Innocent weed can become tangled in wool, and as a result wool is devalued and animals are more difficult to handle. The burrs may also cause ulcers in the mouths of grazing animals.

### **Persistence**

Innocent weed does not establish well in pastures, particularly when there are competitive pasture species present. Burrs lying on the soil surface do not readily germinate, due to low moisture and also inhibiting effect of light, but buried seeds germinate more easily, requiring ongoing control.

### **Policy**

#### **National Strategy**

Innocent weed is not classified as a Weed of National Significance, therefore there is no National Strategy for this species.

#### **State Policy**

To prevent Innocent weed being introduced to areas free of the plant especially urban areas, areas with sandy soils and irrigation areas, under the *Landscape SA Act 2019*:

- Prohibit movement of Innocent weed on public roads.
- Prohibit sale of Innocent weed or produce or goods carrying the plant.
- Require land owners to control Innocent weed on their properties.
- Allow recovery of costs of roadside control of Innocent weed by Landscape Board.

#### **SA Arid Lands Landscape Board Risk Assessment**

The SAAL Landscape Board risk assessment is to **PROTECT SITES** from the spread of Innocent weed.

Aim: To prevent spread of Innocent weed to key sites/assets of high economic, environmental and/or social value.

#### **Port Augusta - Quorn District Risk Management Action**

As above.

### **Best Practice Control**

<b>Control method and description</b>	<b>Best time to control</b>	<b>Link for PIRSA current best practice chemical control and rates</b>
<b>Foliar Spray</b>		
Useful on all plants Spray the entire plant with the recommended herbicide mixture when it is actively growing and before seed set, i.e. within the first week after summer rainfall.	Within 1 week after summer rain	<a href="#">Innocent weed - PIRSA</a>
<b>Physical and/or Organic Control</b>		
Isolated occurrences of Innocent weed can be easily grubbed out of the ground. Grub the plant(s) out of the ground within one week after rain, prior to seed set. Organic foliar spraying can be carried out using Bioweed (680g/L pine oil) at a rate of 20L per 100L water. Complete coverage of plant is required and should be applied when the plant is actively growing. This mixture will also kill seed it contacts both on the plant and on the ground.		

## **Recommended Actions for Innocent weed Management in Port Augusta-Quorn District**

- 1.** Support landholder education in identification, monitoring and control of Innocent weed.
- 2.** Identify current distributions of Innocent weed in neighbouring Landscape regions/districts and ascertain possible pathways of entry.
- 3.** Encourage landholders to undertake control of all Innocent weed infestations in close proximity to key sites.
- 4.** Encourage landholders to monitor success, following control of Innocent weed, and carry out follow up control as necessary.
- 5.** Prevent sale and movement of Innocent weed within the Port Augusta-Quorn District.

## **Surveys and Monitoring**

Keep an eye out for Innocent weed during routine pastoral management activities especially around stock yards and immediately after summer rainfall.

# Khaki weed *Alternanthera pungens*

## Description

Prostrate herb with perennial root system and annual above ground growth. Tap-roots are often large, woody and deep-penetrating. Stems shortly silky hairy. Leaves ovate to circular, hairless to sparsely hairy, margins entire, leaf stalk 0.2–1 cm long. Inflorescence ovoid, 6–10 mm wide, flowers from March to April. Fruit 1–1.5 mm long. Seeds about 1 mm wide, brownish, globe-shaped. It forms carpets of sharp burrs that can injure people and animals. Originally from Brazil, Ecuador, Peru and Venezuela, it was first recorded in NSW in 1989 and first reported in Alice Springs in 1957.

Khaki weed may be confused with other species. For more information click [here](#).



Khaki weed runners (Credit: [www.weeds.org.au](http://www.weeds.org.au))



Khaki weed burrs (Credit: [www.depi.vic.gov.au](http://www.depi.vic.gov.au))

## Distribution

**Australian distribution:** Khaki weed occurs in all mainland states and territories. In New South Wales it occurs in towns, in coastal areas north of Sydney and also agricultural regions in the central and north-west of the state. In Victoria and South Australia, Khaki weed is confined to towns in the more arid areas. It is widespread in Queensland particularly around towns in the south-east. In the Northern Territory, it occurs in the Darwin, Gulf, Katherine, Victoria River and Alice Springs districts. In Western Australia, Khaki weed can be found in and around towns in the Kimberley and Pilbara regions and around Perth and other centres in the south-west.

**SA distribution:** Records of Khaki weed infestations are scattered in the Eyre Peninsula, Northern and Yorke, Kangaroo Island, SA Murray Darling Basin and South East regions, extending as far north as Marla and Vulkathunha – Gammon Ranges National Park in the pastoral zone. Many of these have been eradicated but its distribution is maintained by incursions on vehicles.

**District distribution:** A number of infestations have been reported in Port Augusta.

**Potential distribution:** Khaki weed likes light soils and relies on summer rainfall, so providing these conditions are met, Khaki weed can exist throughout mainland Australia. Due to its spiny seeds, it can easily be picked up on vehicle tyres and spread anywhere vehicles travel.

## Threats and Impacts

### Invasiveness

Spread by seed within spiny bracts that adhere to tyres, clothing and animals. Local spread may also occur through spreading stems that root at nodes. Cultivation can also encourage spread.

### Impacts

Major weed of warm temperate and tropical areas around the world. Stock graze young plants. Widespread in wasteland, caravan parks, orchards and recreation areas. The sharp spiny fruit cause mechanical damage to the feet and mouths of stock and working animals when present in hay and pasture. Khaki Weed is believed to be poisonous to animals and to cause skin ailment in cattle. It is not readily eaten but sometimes young plants are consumed by sheep, apparently without ill effect. It is also claimed to cause hay fever, asthma and dermatitis in some people.

## Persistence

Khaki weed is a difficult plant to control as it is deep-rooted, the tap root also allowing it to survive periods of drought. It also forms a soil seed-bank under infestations, with seed surviving for more than 5 years.

## Policy

### National Strategy

Khaki weed is not classified as a Weed of National Significance, therefore there is no National Strategy for this species.

### State Policy

No losses to the amenity value of recreational land or pasture production due to Khaki weed.

Objectives:

- Detect and destroy all Khaki weed infestations
- Prevent further spread and establishment in SA.

Implementation

- Prohibit entry to area
- Prohibit movement of Khaki weed on public roads
- Prohibit the sale of Khaki weed or contaminated produce
- Require landholders to **notify** Landscape Board of Khaki weed infestations
- Landowners to destroy Khaki weed on their properties
- Recovery of control costs on adjoining road reserves by the Landscape Board

### SA Arid Lands Landscape Board Risk Assessment

SAAL Landscape Board risk assessment for Khaki weed is to **DESTROY INFESTATIONS**.

Aim: To significantly reduce the extent of Khaki weed in the SA Arid Lands Landscape Region through destruction of known infestations.

### Port Augusta-Quorn District Risk Management Action

As above.

### Best Practice Control

Control method and description	Best time to control	Link for PIRSA current best practice chemical control and rates
<b>Foliar Spray</b>		
Spot spraying with a knap sack or quad/ute mounted spray unit can be used. Spray the entire plant with the recommended herbicide mix. Soil residual. Do not use near desirable vegetation or in waterways.	Anytime	<a href="#">Khaki weed - PIRSA</a>
Spot spraying with a knap sack or quad/ute mounted spray unit can be used. Spray the entire plant with the recommended herbicide mix Non-selective, avoid contact with desirable plants. Won't kill the root system, ongoing treatment required.	Whilst plant actively growing	
Fallow. Boom spray. DO NOT treat weeds which are beyond the rosette stage. Use higher rate on large weeds. DO NOT HARVEST, GRAZE OR CUT FOR	Apply when weeds have at least 3-5 true leaves and are actively growing.	

### **Physical and/or Organic Control**

---

Individual plants may be grubbed. Ensure the complete removal of the tuber and root system. Dig beside the root 100mm down or more. Destroy any burrs found.

Organic foliar spraying can be carried out using Bioweed (680g/L pine oil) at a rate of 20L per 100L water. Complete coverage of plant is required and should be applied when the plant is actively growing. This mixture will also kill seed it contacts both on the plant and on the ground.

---

### **Recommended Actions for Khaki weed Management in Port Augusta-Quorn District**

1. Support landholder education in identification, monitoring and control of Khaki weed.
2. Occurrences of Khaki weed to be surveyed, mapped and reported to SAAL Landscape immediately.
3. Identify current distributions of Khaki weed in neighbouring Landscape regions/districts and ascertain possible pathways of entry into Port Augusta-Quorn District.
4. Destroy all infestations as they are detected, aiming for local eradication.
5. Encourage landholders to monitor success following control, and carry out follow up control as necessary.
6. Prevent sale and movement of Khaki weed within Port Augusta-Quorn District.

### **Surveys and Monitoring**

Keep an eye out for Khaki weed during routine pastoral management activities, particularly from spring to autumn. Record locations with GPS or appropriate app and advise SA Arid Lands Landscape Board of infestation.

# Mesquite *Prosopis species*

## Description

Mesquite are a group of shrubs or small trees, which reproduce by seed and suckers. Leaves are fern-like, often with one or two thorns. Flowers are greenish-yellow, small, arranged on cylindrical-shaped flower heads 50-80 mm long, in spring and early summer. Seed pods, up to 20 cm long, are generally straight or slightly curved, and resemble a string of beads. Originally from USA and Mexico, introduced in 1900 as an ornamental plant, fodder and shade tree. There are around 44 species of *Prosopis*, and only a few have been introduced to Australia.

**Do not confuse** Mesquite with prickly acacia trees. For more information click [here](#).



Mature Mesquite tree (Credit: B Shepherd)



Mesquite leaves and thorns (Credit: B Shepherd)



Mesquite flowers (Credit: B Shepherd)

## Distribution

**Australian distribution:** Mesquite occurs in every mainland state. Mesquite has infested large tracks of pastoral country in the semi-arid and arid areas of QLD and WA.

**SA distribution:** Mesquite was infrequently planted in gardens around SA. There are four known naturalised populations of Mesquite in northern SA, in the eastern districts around Cockburn, around Woomera, on the south eastern side of Lake Torrens and another near Port Augusta.

**District distribution:** There is a population of Mesquite in the low sand dunes on the north eastern edge of Port Augusta. The population has persisted despite low average rainfall and sporadic control efforts for over 10 years.

**Potential distribution:** Mesquite's combination of a long life cycle, ability to survive drought, high seed production and long seed life makes it well suited to establishment in northern SA. Mesquite has the ability to establish in the rivers, swamps and other areas where soil moisture persists, throughout northern SA.

## Threats and Impacts

### Invasiveness

The occurrences of Mesquite in SA are on the outer extent of their preferred range and it tends not to spread as rapidly as in the northern Australian climate. However, vigorous establishment and growth of the SA populations demonstrate its potential to successfully invade and establish. Mesquite reproduces from seeds that are readily eaten and dispersed by animals.

### Impacts

Mesquite forms dense thickets that prevent stock from accessing water points and make mustering difficult. The thickets overrun grazing land, exclude all other vegetation and use valuable ground water.

### Persistence

Mesquite will grow in most of the semi-arid and arid environments of northern SA however it prefers areas prone to flooding or where soil moisture is retained. Mesquite is a very hardy plant that can tolerate dry conditions and seeds remain viable for several years.

## Policy

### National Strategy

Mesquite species and hybrids are confined and eventually eradicated from Australia.

### State Policy

To maintain production from semi-arid pastoral lands and protect the integrity of native vegetation, under the *Landscape SA Act 2019*:

- Prohibit entry of Mesquite into SA and movement on public roads.
- Prohibit sale of Mesquite or contaminated material.
- Require landholders to **notify** Landscape Board of Mesquite infestations.
- Require landholders to destroy Mesquite on their properties.
- Recovery of control costs on adjoining road reserves by the Landscape Board.

### SA Arid Lands Landscape Board Risk Assessment

The SAAL Landscape Board risk assessment for Mesquite is to **DESTROY INFESTATIONS**.

Aim: To significantly reduce the extent of Mesquite in the region.

### Port Augusta-Quorn District Risk Management Action

As above.



## Best Practice Control

Control method and description	Best time to control	Link for PIRSA current best practice chemical control and rates
<b>Basal Bark</b>		
<p>Useful for smaller to mid-sized plants that have not developed rough bark.</p> <p>Spray all sides of each stem with the recommended herbicide mix, from the ground up to a height of between 500 - 750mm.</p> <p>For seedlings spray the entire plant.</p>	<p>Most of the year assuming the plant is not stressed. Most effective when actively growing.</p>	<p><a href="#">Mesquite - PIRSA</a></p>
<b>Foliar Spray</b>		
<p>Best for seedlings &amp; regrowth of previously treated areas on plants under 1.5m height</p> <p>Spray the entire plant</p>	<p>Best time to spray is when plants are actively growing and soil moisture is good.</p>	<p><a href="#">Mesquite - PIRSA</a></p>
<b>Cut Stump</b>		
<p>Useful for medium to larger trees</p> <p>The main stem(s) are cut off by chainsaw and the stump immediately (within 10 seconds) painted or sprayed with the recommended herbicide mixture.</p> <p>The stump should be cut as close to the ground as possible.</p>	<p>Any time of year</p>	<p><a href="#">Mesquite - PIRSA</a></p>
<b>Physical and/or Organic Control</b>		
<p>Useful for very large trees, dense infestations and broad scale germinations or regrowth - the entire plant is removed from the ground using a suitably sized bull dozer or loader.</p> <p>Roots must be cut about 30-50 cm below the soil surface to reduce the incidence of regrowth. This is best done with a blade plough.</p> <p>Where regrowth occurs spray using the foliar method.</p> <p>Branch and root fragments of Mesquite can take root, therefore woody material must be moved out of creek lines and moist areas.</p> <p>Ongoing inspections and hand-pulling of seedlings will additionally be required where recruitment is occurring, until the seedbank is exhausted.</p>		

## Recommended Actions for Mesquite Management in Port Augusta-Quorn District

1. Support landholder education in identification, monitoring and control of Mesquite.
2. Encourage landholders to report any infestations of Mesquite they detect so that it can be surveyed, mapped and control sites monitored.

3. Destroy all Mesquite from the Port Augusta-Quorn District including removal of cultivated plants.
4. Encourage landholders to monitor success, following control of Mesquite, and carry out follow up control as necessary.
5. Consider quarantine provisions around known occurrences of Mesquite.
6. Prevent sale and movement of Mesquite within SAAL Landscape region.

### **Surveys and Monitoring**

Keep an eye out for Mesquite during routine pastoral management activities. Record locations with GPS or appropriate app and advise SA Arid Lands Landscape Board of infestation. Monitor any time of year, look for fern like leaves, long yellow flower heads and sometimes bright light green leaves.

# Parkinsonia *Parkinsonia aculeata*

## Description

Parkinsonia is a much branched shrub or tree, 2-8 m tall but at times reaching 10 m tall. Parkinsonia has extensive surface roots and a deep taproot. Leaves consist of a flat, green leaf stalk (up to 30 cm long) with numerous pairs of small oval leaflets, often with a sharp spine (3-20 mm long) Flowers are fragrant up to 2 cm diameter, bright yellow in colour becoming reddish with age, summer to early autumn. The top petal either has orange spots or turns completely orange. The pod is almost straight (3-13 cm long and 3.5-8 mm wide), narrowing at each end.

Originally from Central America, Parkinsonia was introduced in the 1890s to early 1900s as a shade and ornamental shrub. Parkinsonia has the potential to invade more than three quarters of mainland Australia.

**Do not confuse** Mesquite with prickly acacia trees. For more information click [here](#).



Parkinsonia tree (Credit: B Shepherd)



Parkinsonia flowers (Credit: B Shepherd)



Parkinsonia seed pods (Credit: B Shepherd)

## Distribution

**Australian distribution:** Parkinsonia was commonly planted in many areas around Australia in the late 1800s. Over one million hectares of Australia is now infested with Parkinsonia primarily throughout coastal, central and western QLD, central and northern NT and the Kimberly and Pilbara regions of WA.

**SA distribution:** Within northern SA there are 3 current locations where Parkinsonia has naturalised – near Leigh creek in northern Flinders Ranges, around Woomera and Port Augusta. All these occurrences are small populations (2-10 plants). Historically there have been more occurrences recorded within gardens, however all of these have been successfully controlled.

**District distribution:** There are scattered Parkinsonia infestations in the Port Augusta area.

**Potential distribution:** Parkinsonia has the potential to establish in watercourses, swampy areas and other watered areas throughout arid and semi-arid areas of northern SA.

## Threats and Impacts

### Invasiveness

Parkinsonia reproduces by seeds that are spread by flood waters and in mud attached to vehicles or animals. It establishes and grows in a wide variety of conditions from semi-arid to sub-humid environments and prefers drainage lines, swamps, floodplains and other watered areas. Mature plants typically produce 5000 seeds annually, and germination after flood event or disturbance can be prolific.

### Impacts

Parkinsonia forms dense thickets and can take over vast tracks of land replacing native plant species including valuable pasture species. It reduces access to land and water and increases difficulty when mustering stock. Parkinsonia reduces habitat for native species and provides a refuge for feral animals. The areas at risk are the more productive arid rivers and floodplains.

### Persistence

Parkinsonia can re-shoot from roots. Germination from seed after control disturbance can be prolific. Parkinsonia seeds remain viable for many years.

## Policy

### National Strategy

Parkinsonia is confined and its impact reduced to a minimum.

### State Policy

To maintain access to pastoral lands, and the integrity of native vegetation, under the *Landscape SA Act 2019*:

- Prohibit movement of Parkinsonia into SA.
- Prohibit movement of Parkinsonia on public roads.
- Prohibit sale of Parkinsonia, or produce or goods carrying Parkinsonia.
- Require landholders to **notify** Landscape Board of Parkinsonia infestations.
- Require landholders to destroy Parkinsonia on their properties.
- Recovery of control costs on adjoining road reserves by the Landscape Board.

### SA Arid Lands Landscape Board Risk Assessment

The SAAL Landscape Board risk assessment for Parkinsonia is to **DESTROY INFESTATIONS**.

Aim: To significantly reduce the extent of Parkinsonia in the region.

### Port Augusta-Quorn District Risk Management Action

As above

### Best Practice Control

Control method and description	Best time to control	Link for PIRSA current best practice chemical control and rates
<b>Basal Bark</b>		
Useful for smaller to mid-sized plants that have not developed rough bark.	Most of the year assuming	<a href="#">Parkinsonia - PIRSA</a>

---

Spray all sides of each stem with the recommended herbicide mix, from the ground up to a height of between 500 - 750mm. For seedlings spray the entire plant.	the plant is not stressed. Most effective when actively growing.
--	--

---

### **Foliar Spray**

---

Best for seedlings & regrowth of previously treated areas on plants under 1.5m height Spray the entire plant	Best time to spray is when plants are actively growing and soil moisture is good.	<a href="#">Parkinsonia - PIRSA</a>
---	---	-------------------------------------

---

### **Cut Stump**

---

Useful for medium to larger trees The main stem(s) are cut off by chainsaw and the stump immediately (within 10 seconds) painted or sprayed with the recommended herbicide mixture. The stump should be cut as close to the ground as possible.	Any time of year	<a href="#">Parkinsonia - PIRSA</a>
--	------------------	-------------------------------------

---

### **Biological Control**

---

Drill and insert 1 or 2 capsules into tree and seal with dowel. Contact Community Landscape Officer for more information.	Any time of year	<a href="#">Parkinsonia - PIRSA</a>
---	------------------	-------------------------------------

---

### **Physical Control**

---

Useful for very large trees, dense infestations and broad scale germination or regrowth.  
The entire plant is removed from the ground using a suitable bulldozer or loader.  
Roots must be cut about 30-50 cm below the soil surface to reduce the incidence of regrowth. This is best done with a blade plough.  
Branch and root fragments of Parkinsonia can take root, therefore woody material must be moved out of creek lines and moist areas.  
Ongoing inspections and hand-pulling of seedlings will additionally be required, where recruitment is occurring, until the seedbank is exhausted.

---

## **Recommended Actions for Parkinsonia Management in Port Augusta-Quorn District**

1. Support landholder education in identification, monitoring and control of Parkinsonia.
2. Encourage landholders to provide data on distribution of Parkinsonia so that it can be surveyed and mapped.
3. Destroy all infestations as they are identified, aiming for local eradication.
4. Encourage landholders to monitor success, following control of Parkinsonia, and carry out follow up control as necessary.
5. Prohibit sale and movement of Parkinsonia within Port Augusta-Quorn District.

## **Surveys and Monitoring**

Keep an eye out for Parkinsonia during routine pastoral management activities. Record locations with GPS or appropriate app and advise SA Arid Lands Landscape Board of infestation. Monitor any time of the year, look for the distinctive small leaves and the overall bright light green colour of the plant.

## Appendix 1 – Noted Weeds

The weeds in this section are not declared under the *Landscape SA Act 2019* and therefore are not considered priority weeds by the Port Augusta – Quorn Landscape Group. However, they are included in this appendix for identification for local landholders and other stakeholders.

### Castor Oil Plant *Ricinus communis*

#### Description

Tall, branching perennial shrub which grows to 3 m high and occasionally higher. It has stout hollow branches which are a dull pale green or red, older branches and trunks turn greyish. Large leaves (10–60 cm across) are widely spaced on the branches. Each leaf is divided into 7–9 pointed triangular segments with toothed edges and conspicuous veins. Leaves are glossy, dark reddish-green when young, glossy green when mature. Flowers are crowded in stout, erect spikes in the forks of the upper branches, from summer to autumn. Female flowers are in the upper part of the spikes, male flowers at the base. Female flowers develop into fruits about 2.5 cm across which are covered with soft green or red spines. These fruits have three segments, each segment containing one large, mottled, smooth seed. When ripe, the fruits explode violently and throw the seeds a distance of several metres. Originally from Africa and Eurasia it was first recorded in Australia in 1803 and was introduced as an ornamental for gardens. The seed contains ricin and is poisonous to stock and people. The sap can cause temporary blindness if exposed to eyes.



Castor Oil Plant (Credit: P Hodges)



Castor Oil Plant with fruit (Credit: DAFF, QLD)

#### Distribution

**Australian distribution:** Found in all mainland States of Australia.

**SA distribution:** Found in Mt Lofty Ranges, Fleurieu Peninsula, South East, Flinders Ranges, North East Pastoral, Mid North and Murray Mallee.

**District distribution:** Found in some drainage lines and disturbed roadsides in the District.

**Potential distribution:** Mainly in drainage lines but spreads slowly. It has been naturalised in areas of Australia for a long time and its current extent is not anticipated to greatly change.

## Threats and Impacts

### Invasiveness

Invasive in disturbed areas. Spread locally several metres by the explosive release of seed from the ripening capsule. Water flow, road maintenance and mud on vehicles also contribute to its spread. Large plants only develop in wet areas such as creeks, where they may cause blockages. Generally spreads slowly.

### Impacts

The seeds of castor oil contain ricin, a poison that is extremely toxic to livestock and humans. Leaves have a lesser amount of toxin. Symptoms of poisoning in animals usually do not appear for a few hours or several days. However, stock find castor oil unpalatable and won't eat it if alternative feed is available.

### Persistence

Seeds may emerge from 30cm below the soil surface. Seed has medium longevity. Drought tolerant. Very rapid growth rates and may seed in its first year.

## Policy

### National Strategy

Castor Oil Plant is not classified as a Weed of National Significance, therefore there is no National Strategy for this species.

### State Policy

Castor Oil Plant is not a declared species under the *Landscape SA Act 2019*, therefore there is no South Australian policy.

### SA Arid Lands Landscape Board Risk Assessment

The SAAL Landscape Board risk assessment for Castor Oil Plant is **LIMITED ACTION**.

Aim: Castor Oil Plant would only be targeted for coordinated control in the management area if its local presence makes it likely to spread to land uses where it ranks as a higher priority.

### Port Augusta - Quorn District Risk Management Action

As above.

## Best Practice Control

Control method and description	Best time to control	Link for PIRSA current best practice chemical control and rates
<b>Foliar Spray</b>		
Spot spray.	While plant is actively growing	None listed
<b>Physical</b>		
Seedling plants may be removed by hand-pulling. Use protective gloves and glasses to prevent sap contacting skin or eyes.		
Larger plants can be mechanically removed using a loader. Burn or dispose of the plants as preferred. Soil disturbance associated with this method will provide ideal conditions for seed germination and follow-up control will be necessary.		



## Recommended Actions for Castor Oil Plant Management in Port Augusta - Quorn District

1. Support landholder education in identification, monitoring and control of Castor Oil Plant.
2. Encourage landholders to provide data on distribution of Castor Oil Plant so that it can be mapped and surveyed.
3. Provide input into identification of key sites for control of Castor Oil Plant infestations.
4. Assist in identifying and coordinating Castor Oil Plant control programs using integrated weed management, especially in areas where key sites are threatened.
5. Encourage landholders to monitor success, following control of Castor Oil Plant, and carry out follow up control as necessary.

### Surveys and Monitoring

Keep an eye out for Castor Oil Plant during routine pastoral management activities, particularly along drainage lines and record locations using a GPS.

## Mexican poppy *Argemone ochroleuca*

### Description

Stiff bluish-green prickly plant that grows up to 1 m in height. Leaves up to 20cm long, silvery green with white veining and deep regular lobes. The upper surface of the leaf is smooth while the underside has a few prickles along the midrib. The stem leaves, in contrast to the rosette leaves, are stalkless and clasp onto the plant's stem. Flowers 6 cm in diameter with four light yellow or cream petals, flowering generally November to February but is opportunistic in arid areas and can flower whenever moisture is available. Fruit spiny, oblong seed capsules, 3.5cm in length with three to six openings at the top. The capsule contains up to 400 seeds. Seeds are oval-shaped, blackish brown, about 1.5 mm long with a pitted seed coat and a ridge along one side. Originally from Mexico, Central America and USA, it was first reported near Sydney in 1845, probably imported as a contaminant of wheat seed. *Argemone ochroleuca* is similar in appearance and far more widespread than *A. mexicana*, and both species are referred to as Mexican poppy. *A. mexicana* differs from *A. ochroleuca* in that it has bright yellow flowers as opposed to cream or pale-yellow flowers, and globular flower buds as opposed to the egg-shaped buds of *A. ochroleuca*.



Seed pod and flower (Credit: DAFF)



Flower and spiny leaves (Credit: DAFF)

### Distribution

**Australian distribution:** All Australian states and territories, except Tasmania.

**SA distribution:** Some intermittent creek-lines, roadsides and areas of soil disturbance within the SA Arid Lands and Northern and Yorke regions.

**District distribution:** Isolated patches in creek-lines around Hawker. Found in intermittent creek-lines and areas of disturbed soil.

**Potential distribution:** Potential distribution of Mexican poppy extends across the range lands mainly in drainage lines and along roadsides.

## Threats and Impacts

### Invasiveness

Establishes readily on disturbed ground, overgrazed pastures, creek beds and roadsides. Seed is readily dispersed by moving water, especially where plants grow in riparian habitats and on terrain where erosion and runoff occurs. Contaminated soil, fodder, vehicles and the fur and hooves of livestock are vectors for transport of Mexican poppy seed.

### Impacts

The prickly fruits can become entangled in wool, reducing its value considerably. It is suspected of being toxic to animals and humans due to the presence of alkaloids in all parts of the plant. However, reported cases of poisoning are rare because livestock tend to avoid it in the field, due to the presence of a bitter sap that makes it unpalatable. Poisonings may occur as a result of hay and chaff containing traces of the plant. Poisonings in humans have occurred by the consumption of oils contaminated with the plant.

### Persistence

A single plant can produce up to 30 000 seeds per year. The seed can stay dormant for many years, making control difficult.

## Policy

### National Strategy

Mexican poppy is not classified as a Weed of National Significance, therefore there is no national strategy for this species.

### State Policy

Mexican poppy is not a declared species under the *Landscape SA Act 2019*, therefore there is no South Australian policy.

### SA Arid Lands Landscape Board Risk Assessment

SAAL Landscape Board risk assessment for Mexican poppy is to **MONITOR** for any changes in the species weed risk.

Aim: To detect any significant changes in Mexican poppy's weed risk and monitor the spread of the species and review any changes in weediness.

### Port Augusta-Quorn District Risk Management Action

As above.

## Best Practice Control

---

### Foliar Spray

---

While no herbicides are registered specifically for this weed, in general, it's legal to use a registered herbicide in accordance with the label directions, i.e. method of application and situation, on any weed, not just the ones mentioned by name on the manufacturer's label.

---

### Physical and/or Organic Control

---

---

Hand pulling or grubbing using a mattock/hoe for immature plants that haven't seeded.

If you locate a plant with mature seeds, apart from removing plant, collect seeds and burn them (e.g. campfire).

---

### **Recommended Actions for Mexican poppy Management in Port Augusta-Quorn District**

1. Support landholder education in identification, monitoring and control of Mexican poppy.
2. Monitor distributions of Mexican poppy over time through the use of surveillance and mapping.
3. Provide input into identification of key sites requiring control of Mexican poppy.
4. Encourage landholders to monitor success, following control and carry out follow up control as necessary.

### **Surveys and Monitoring**

Monitor during routine pastoral management activities, particularly late spring to late autumn and after summer rains, and especially along roadsides and drainage lines.

# Pepper tree *Schinus molle*

## Description

A large evergreen tree to 12 m tall and 5-10 m wide. Leaves are divided into 15-41 alternatively arranged, shiny leaflets, 4 cm long. Leaves are aromatic and sticky when crushed. Small white flowers hang in long sprays, all year round, and are followed by spherical shiny pink fruit, 4-6 mm in diameter. Originally from South America, it was introduced in 1870s as a shade tree. All *Schinus* species leaves and fruit are poisonous to livestock and possibly humans.



Mature Pepper tree (Credit: P Hodges)



Immature fruit (Credit: B Shepherd)



Mature fruit (Credit: B Shepherd)

## Distribution

**Australian distribution:** Pepper trees are favoured garden plants throughout Australia and their ability to withstand dry conditions makes them well suited as garden and shade plantings in arid and semi-arid areas.

**SA distribution:** Common at nearly every township, station house and stockyard in northern SA. Naturalised Pepper trees are a common site, particularly along drainage lines of the Flinders Ranges, the Olary Ranges and in the mid north agricultural districts. Within some creeks and swamps, naturalised Pepper trees are the dominant tree choking out all other plants.

**District distribution:** Pepper trees were planted at most stations throughout the Port Augusta - Quorn District as amenity trees. There are numerous locations where Pepper trees have naturalised along adjacent creek lines across the District.

**Potential distribution:** Pepper trees have demonstrated their ability to establish in many of the semi-arid drainage lines and swampy areas of SA.

## Threats and Impacts

### Invasiveness

Pepper trees reproduce from seed, and germination requires sustained soil moisture. The seeds are dispersed by birds, animals, water and human activity. Broad scale seed germinations can occur following floods and favourable rainfall events, especially in swampy areas.

### Impacts

Pepper trees can form dense thickets in swamps and along rivers excluding native vegetation and using valuable ground water.

### Persistence

Mature Pepper trees can re-shoot following initial chemical control and can re-shoot from root portions left in the ground following mechanical control. Seeds remain viable for less than three years.

## Policy

### National Strategy

Pepper trees are not classified as a Weed of National Significance, therefore there is no National Strategy for this species.

### State Policy

Pepper tree is not a declared species under the *Landscape SA Act 2019*, therefore there is no South Australian policy.

### SA Arid Lands Landscape Board Risk Assessment

The SAAL Landscape Board risk assessment is to **PROTECT SITES** from Pepper tree infestation. Aim: To prevent spread of Pepper trees to key sites/assets of high economic, environmental and/or social value, through targeted control.

### Port Augusta - Quorn District Risk Management Action

As above.

## Best Practice Control

---

### Cut Stump

---

While no herbicides are register specifically for this weed, in general, it's legal to use a registered herbicide in accordance with the label directions, ie method of application and situation, on any weed, not just the ones mentioned by name on the manufacturer's label.

---

### Basal Bark

---

While no herbicides are registered specifically for this weed, in general, it's legal to use a registered herbicide in accordance with the label directions, ie method of application and situation, on any weed, not just the ones mentioned by name on the manufacturer's label.

---

---

### **Physical and/or Organic Control**

---

Useful for very large trees, dense infestations and broad scale germinations or regrowth - the entire plant is removed from the ground using a suitably sized bull dozer or loader.

Roots must be cut about 30-50 cm below the soil surface to reduce the incidence of regrowth. This is best done with a blade plough.

Where regrowth occurs spray using a foliar method.

Best time to use this method - Summer or when plants are stressed e.g. during drought conditions.

---

### **Recommended Actions for Pepper tree Management in Port Augusta - Quorn District**

- 1.** Support landholder education in identification, monitoring and control of Pepper trees.
- 2.** Provide input into identification of high priority areas such as significant rivers, creeks and waterholes and coordinate control programs to protect identified priority sites.
- 3.** Encourage landholders to control naturalised Pepper trees, and control cultivated Pepper trees where key sites are threatened.
- 4.** Encourage landholders to monitor success, following control of Pepper trees, and carry out follow up control as necessary.

### **Surveys and Monitoring**

Keep an eye out for naturalised Pepper trees during routine pastoral management activities. Monitor any time of the year, look for the distinctive bi-pinnate leaves and the overall light green colour of the plant.

## Appendix 2 – Cactus and appropriate biocontrol species

Species name	Common name/s	Biological control agent/s	Effectiveness of agent
<b>Species for which agents provide effective control</b>			
<i>Cylindropuntia fulgida</i> var. <i>mamillata</i>	Boxing glove cactus, coral cactus	<i>Dactylopius tomentosus</i> 'cholla' lineage	Effective control provided.
<i>Cylindropuntia imbricata</i>	Rope pear, devil's rope pear	<i>Dactylopius tomentosus</i> 'cylindropuntia' lineage	Effective control provided. Released at several sites in north-western and western NSW, where field observations indicate this lineage is more damaging than 'imbricata'.
			Released at a small number of sites in Qld.
		<i>Dactylopius tomentosus</i> 'imbricata' lineage	Effective control provided. Prevalent in Qld and SA.
		<i>Dactylopius tomentosus</i> 'bigelovii' lineage	Research indicates good potential however further testing required.
<i>Cylindropuntia kleiniiae</i>	Klein's cholla	<i>Dactylopius tomentosus</i> 'imbricata' lineage	Effective control provided.
		<i>Dactylopius tomentosus</i> 'bigelovii' lineage	Research indicates good potential however further testing required.
<i>Cylindropuntia leptocaulis</i>	Candle cholla	<i>Dactylopius tomentosus</i> 'imbricata' lineage	Effective control provided.
<i>Cylindropuntia pallida</i>	White spined Hudson pear	<i>Dactylopius tomentosus</i> 'californica var. parkerii' lineage	Effective control provided in some locations, but has not proven to be effective in SA as yet.
<i>Cylindropuntia spinosior</i>	Snake cactus	<i>Dactylopius tomentosus</i> 'bigelovii' lineage	Effective control provided. Released at a small number of sites in NSW where it is impactful once established (which can take time).
		<i>Dactylopius tomentosus</i> 'spinosior-safford' lineage	Research indicates good potential however further testing required.
<i>Cylindropuntia prolifera</i>	Jumping cholla	<i>Dactylopius tomentosus</i> 'californica var. parkerii' lineage	Effective control provided.
<i>Cylindropuntia tunicata</i>	Brown spined Hudson pear	<i>Dactylopius tomentosus</i> 'acanthocarpa var. echinocarpa' lineage	Effective control provided.
<i>Opuntia aurantiaca</i>	Tiger pear	<i>Dactylopius austrinus</i>	Effective control provided.
		<i>Cactoblastis cactorum</i>	Provide some control, though not as effective as <i>Dactylopius austrinus</i> .
		<i>Tucumania tapiacola</i>	

<i>Opuntia elata</i>	Riverina pear	<i>Cactoblastis cactorum</i>	Effective control provided.
		<i>Dactylopius opuntiae</i> 'ficus' lineage	Further testing of <i>Dactylopius</i> in progress to determine effectiveness.
		<i>Dactylopius ceylonicus</i>	
<i>Opuntia englemannii</i>	Engelmann's prickly pear	<i>Dactylopius opuntiae</i> 'ficus' lineage	Effective control provided.
<i>Opuntia monacantha</i>	Smooth tree pear	<i>Dactylopius ceylonicus</i>	Effective control provided. Takes several years to kill plants.
		<i>Cactoblastis cactorum</i>	Attacks plants and limits growth but does not control it.
<i>Opuntia robusta</i>	Wheel cactus	<i>Dactylopius opuntiae</i> 'ficus' lineage	Effective control provided.
<i>Opuntia stricta</i>	Common prickly pear	<i>Cactoblastis cactorum</i>	Effective control provided.
		<i>Dactylopius opuntiae</i> 'stricta' lineage	
<i>Opuntia tomentosa</i>	Velvety tree pear	<i>Cactoblastis cactorum</i>	Effective control provided on young, small plants.
		<i>Dactylopius opuntiae</i> 'stricta' lineage	Effective control provided.
<b>Further work required to identity an agent or to determine agent effectiveness</b>			
<i>Opuntia elatior</i>	Red-flower prickly pear	<i>Dactylopius opuntiae</i> 'ficus' lineage	Effective control provided in NSW based on field observations. Released in SA in 2022, however field efficacy has not been evaluated.
<i>Opuntia humifusa</i>	-	<i>Dactylopius opuntiae</i> 'stricta' lineage	South African research indicates this lineage may be suitable, however there is a lack of data in Australia to confirm.
<i>Opuntia puberula</i>	-	None currently available.	Further testing of <i>Dactylopius</i> required
<i>Opuntia schickendantzii</i>	Chicken dance cactus	None currently available.	Further testing required to confirm efficacy
<i>Opuntia streptacantha</i>	Westwood pear	None currently available.	Further testing of <i>Dactylopius</i> required

Credit: Shauna Potter, Dept. of Primary Industries and Regional Development, W.A.



## New Infestations or Unknown Weeds

If you find a new infestation of a weed you haven't had before or a weed species that you don't know, the Landscape Board would like to know about it. You can contact the SA Arid Lands Landscape Board on (08) 8429 9666 and ask for the Community Landscape Officer for your District or the Biosecurity Officer.

Please provide the following information when you call - this will help us assist you:

Your name	
Property name	
Paddock name	
GPS location or description	
Description of weed	
Extend of weed	
Description of site where weed is located	

It would also help if you could take some photos of the weed. The Officer you speak to will be able to provide an email address for the photos to be sent to. We will then attempt to identify the weed and we may forward the information and photo to the Weeds Botanist at the State Herbarium.

Published by the SA Arid Lands Landscape Board.  
November 2023

T: +61 (8) 8429 9666

E: SAAL.LandscapeBoard@sa.gov.au



With the exception of the Piping Shrike emblem, other material or devices protected by Aboriginal rights or a trademark, and subject to review by the Government of South Australia at all times, the content of this document is licensed under the Creative Commons Attribution 4.0 Licence. All other rights are reserved.